

NAME _____ DATE _____ PERIOD _____

12-1 Skills Practice

The Counting Principle

State whether the events are *independent* or *dependent*.

1. finishing in first, second, or third place in a ten-person race **dependent**
2. choosing a pizza size and a topping for the pizza **independent**
3. Seventy-five raffle tickets are placed in a jar. Three tickets are then selected, one after the other, without replacing a ticket after it is chosen. **dependent**
4. The 232 members of the freshman class all vote by secret ballot for the class representative to the Student Senate. **independent**

Solve each problem.

5. A surveying firm plans to buy a color printer for printing its maps. It has narrowed its choice to one of three models. Each of the models is available with either 32 megabytes of random access memory (RAM), 64 megabytes of RAM, or 128 megabytes of RAM. From how many combinations of models and RAM does the firm have to choose? **9**
6. How many arrangements of three letters can be formed from the letters of the word *MATH* if any letter will not be used more than once? **24**
7. Allan is playing the role of Oliver in his school's production of *Oliver Twist*. The wardrobe crew has presented Allan with 5 pairs of pants and 4 shirts that he can wear. From how many possible costumes consisting of a pair of pants and a shirt does Allan have to choose? **20**
8. The 10-member steering committee that is preparing a study of the public transportation needs of its town will select a chairperson, vice-chairperson, and secretary from the committee. No person can serve in more than one position. In how many ways can the three positions be filled? **720**
9. Jeanine has decided to buy a pickup truck. Her choices include either a V-6 engine or a V-8 engine, a standard cab or an extended cab, and 2-wheel drive or 4-wheel drive. How many possible models does she have to choose from? **8**
10. A mail-order company that sells gardening tools offers rakes in two different lengths. Customers can also choose either a wooden, plastic, or fiberglass handle for the rake. How many different kinds of rakes can a customer buy? **6**
11. A Mexican restaurant offers chicken, beef, or vegetarian fajitas wrapped with either corn or flour tortillas and topped with either mild, medium, or hot salsa. How many different choices of fajitas does a customer have? **18**

Chapter 12

Glencoe Algebra 2

8

NAME _____ DATE _____ PERIOD _____

12-1 Practice

The Counting Principle

State whether the events are *independent* or *dependent*.

1. choosing an ice cream flavor and choosing a topping for the ice cream **independent**
2. choosing an offensive player of the game and a defensive player of the game in a professional football game **independent**
3. From 15 entries in an art contest, a camp counselor chooses first, second, and third place winners. **dependent**
4. Jillian is selecting two more courses for her block schedule next semester. She must select one of three morning history classes and one of two afternoon math classes. **independent**

Solve each problem.

5. A briefcase lock has 3 rotating cylinders, each containing 10 digits. How many numerical codes are possible? **1000**
6. A golf club manufacturer makes irons with 7 different shaft lengths, 3 different grips, 5 different lies, and 2 different club head materials. How many different combinations are offered? **210**
7. There are five different routes that a commuter can take from her home to the office. In how many ways can she make a round trip if she uses a different route coming than going? **20**
8. In how many ways can the four call letters of a radio station be arranged if the first letter must be W or K and no letters repeat? **27,600**
9. How many 7-digit phone numbers can be formed if the first digit cannot be 0 or 1, and any digit can be repeated? **8,000,000**
10. How many 7-digit phone numbers can be formed if the first digit cannot be 0, and any digit can be repeated? **9,000,000**
11. How many 7-digit phone numbers can be formed if the first digit cannot be 0 or 1, and no digit can be repeated? **483,840**
12. How many 7-digit phone numbers can be formed if the first digit cannot be 0, and no digit can be repeated? **544,320**
13. How many 6-character passwords can be formed if the first character is a digit and the remaining 5 characters are letters that can be repeated? **118,813,760**
14. How many 6-character passwords can be formed if the first and last characters are digits and the remaining characters are letters? Assume that any character can be repeated. **45,697,600**

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.


Chapter 12

Glencoe Algebra 2

9

12-1**Word Problem Practice****The Counting Principle**

1. **CANDY** Amy, Bruce, and Carol can choose one piece of candy from either a white or black bag. The white bag contains various chocolates. The black bag contains small bags of jelly beans. Amy picks from the white bag, and Bruce and Carol both pick from the black bag. Describe whether each of the picks is related as dependent or independent events.



Amy's pick is independent of each of Bruce and Carol's picks; Bruce and Carol's picks are examples of dependent events.

2. **PHOTOS** Morgan has three pictures that she would like to display side by side.



In how many different ways can the pictures be displayed?
6

3. **COMBINATION LOCKS** Eric uses a combination lock for his locker. The lock uses a three number secret code. Each number ranges from 1 to 35, inclusive. How many different combinations are possible with Eric's lock?
42,875

4. **TRUE OR FALSE** Faith is preparing a true or false quiz for her biology class. How many different answer keys can there be for a 10 question true or false quiz?
1024


WEBSITES For Exercises 5-8, use the following information.

Greg is registering to use a website. The website requires him to choose an 8-character alphanumeric password that is not case-sensitive. In other words, for each character, he can choose one of the 26 letters A through Z or one of the 10 digits 0 through 9.


5. How many different passwords are possible?
2,821,109,907,456
6. Greg decides to use a password that does not contain any repeated characters. How many different passwords are possible with this constraint?
1,220,096,908,800
7. Suppose Greg chooses to use only letters with possible repeats. How many different passwords would be possible?
208,827,064,576
8. If Greg's password begins with his first name and ends with his birth month and date, how many possibilities would need to be checked to find his password?
372

12-1**Enrichment****Tree Diagrams and the Power Rule**

If you flip a coin once, there are two possible outcomes: heads showing (*H*) or tails showing (*T*). The tree diagram to the right shows the four (2^2) possible outcomes if you flip a coin twice.



Example 1 Draw a tree diagram to show all the possible outcomes for flipping a coin three times. List the outcomes.




There are eight (2^3) possible outcomes. With each extra flip, the number of outcomes doubles. With 4 flips, there would be sixteen (2^4) outcomes.

The Power Rule for the number of outcomes states that if an experiment is repeated n times, and if there are b possible outcomes each time, there are b^n total possible outcomes.

Find the total number of possible outcomes for each experiment. Use tree diagrams to help you.

- flipping a coin 5 times **2⁵**
- flipping a coin 8 times **2⁸**
- rolling a 6-sided die 3 times **6³**
- rolling a 4-sided die 3 times **4³**
- doing the marble experiment 6 times **3⁶**
- rolling a 6-sided die 2 times **6²**
- rolling a 4-sided die 2 times **4²**
- rolling a 12-sided die 2 times **12²**

Example 2 In a cup there are a red, a blue, and a yellow marble. How many possible outcomes are there if you draw one marble at random, replace it, and then draw another?



There are nine (3^2) possible outcomes.

NAME _____ DATE _____ PERIOD _____

12-2 Lesson Reading Guide

Permutations and Combinations

Get Ready for the Lesson

Read the introduction to Lesson 12-2 in your textbook.

Suppose that 20 students enter a math contest. In how many ways can first, second, and third places be awarded? (Write your answer as a product. Do not calculate the product.)
20 · 19 · 18

Read the Lesson

- Indicate whether each situation involves a *permutation* or a *combination*.
 - choosing five students from a class to work on a special project **combination**
 - arranging five pictures in a row on a wall **permutation**
 - drawing a hand of 13 cards from a 52-card deck **combination**
 - arranging the letters of the word *algebra* **permutation**
- Write an expression that can be used to calculate each of the following.
 - number of combinations of n distinct objects taken r at a time $\frac{n!}{(n-r)!r!}$
 - number of permutations of n objects of which p are alike and q are alike $\frac{n!}{p!q!}$
 - number of permutations of n distinct objects taken r at a time $\frac{n!}{(n-r)!}$
- Five cards are drawn from a standard deck of cards. Suppose you are asked to determine how many possible hands consist of one heart, two diamonds, and two spades.
 - Which of the following would you use to solve this problem: *Fundamental Counting Principle*, *permutations*, or *combinations*? (More than one of these may apply.)
Fundamental Counting Principle, combinations
 - Write an expression that involves the notation $P(n, r)$ and/or $C(n, r)$ that you would use to solve this problem. (Do not do any calculations.)
 $C(13, 1) \cdot C(13, 2) \cdot C(13, 2)$

Remember What You Learned

- Many students have trouble knowing when to use permutations and when to use combinations to solve counting problems. How can the idea of *order* help you to remember the difference between permutations and combinations?
Sample answer: A permutation is an arrangement of objects in which order is important. A combination is a selection of objects in which order is not important.

NAME _____ DATE _____ PERIOD _____

12-2 Study Guide and Intervention

Permutations and Combinations

Permutations When a group of objects or people are arranged in a certain order, the arrangement is called a **permutation**.

Permutations	The number of permutations of n distinct objects taken r at a time is given by $P(n, r) = \frac{n!}{(n-r)!}$.
Permutations with Repetitions	The number of permutations of n objects of which p are alike and q are alike is $\frac{n!}{p!q!}$.

The rule for permutations with repetitions can be extended to any number of objects that are repeated.

Example From a list of 20 books, each student must choose 4 books for book reports. The first report is a traditional book report, the second a poster, the third a newspaper interview with one of the characters, and the fourth a timeline of the plot. How many different orderings of books can be chosen?

Since each book report has a different format, order is important. You must find the number of permutations of 20 objects taken 4 at a time.

$$\begin{aligned}
 P(n, r) &= \frac{n!}{(n-r)!} && \text{Permutation formula} \\
 P(20, 4) &= \frac{20!}{(20-4)!} && n = 20, r = 4 \\
 &= \frac{20!}{16!} && \text{Simplify} \\
 &= \frac{20 \cdot 19 \cdot 18 \cdot 17 \cdot 16 \cdot 15 \cdot \dots \cdot 1}{16 \cdot 15 \cdot \dots \cdot 1} && \text{Divide by common factors.} \\
 &= 116,280
 \end{aligned}$$

Books for the book reports can be chosen 116,280 ways.

Exercises

Evaluate each expression.

- $P(6, 3)$ **120**
- $P(8, 5)$ **6720**
- $P(9, 4)$ **3024**
- $P(11, 6)$ **332,640**
- MOM **3**
- MONDAY **720**
- STEREO **360**

How many different ways can the letters of each word be arranged?

- SCHOOL** The high school chorus has been practicing 12 songs, but there is time for only 5 of them at the spring concert. How many different orderings of 5 songs are possible?
95,040

NAME _____ DATE _____ PERIOD _____

12-2 Skills Practice

Permutations and Combinations

Evaluate each expression.

1. $P(6, 3)$ **120**
2. $P(8, 2)$ **56**
3. $P(2, 1)$ **2**
4. $P(3, 2)$ **6**
5. $P(10, 4)$ **5040**
6. $P(5, 5)$ **120**
7. $C(2, 2)$ **1**
8. $C(5, 3)$ **10**
9. $C(4, 1)$ **4**
10. $C(8, 7)$ **8**
11. $C(3, 2)$ **3**
12. $C(7, 4)$ **35**

Determine whether each situation involves a **permutation** or a **combination**. Then find the number of possibilities.

13. seating 8 students in 8 seats in the front row of the school auditorium
permutation; 40,320
14. introducing the 5 starting players on the Woodsville High School basketball team at the beginning of the next basketball game
permutation; 120
15. checking out 3 library books from a list of 8 books for a research paper
combination; 56
16. choosing 2 movies to rent from 5 movies
combination; 10
17. the first-, second-, and third-place finishers in a race with 10 contestants
permutation; 720
18. selecting 4 candidates to a municipal planning board from a field of 7 candidates
combination; 35
19. choosing 2 vegetables from a menu that offers 6 vegetable choices
combination; 15
20. an arrangement of the letters in the word *rhombus*
permutation; 5040
21. selecting 2 of 8 choices of orange juice at a store
combination; 28
22. placing a red rose bush, a yellow rose bush, a white rose bush, and a pink rose bush in a row in a planter
permutation; 24
23. selecting 2 of 9 kittens at an animal rescue shelter
combination; 36
24. an arrangement of the letters in the word *isosceles*
permutation; 30,240

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____ DATE _____ PERIOD _____

12-2 Study Guide and Intervention (continued)

Permutations and Combinations

Combinations An arrangement or selection of objects in which order is *not* important is called a combination.

The number of combinations of n distinct objects taken r at a time is given by $C(n, r) = \frac{n!}{(n-r)!r!}$.

Example 1 **SCHOOL** How many groups of 4 students can be selected from a class of 20?
Since the order of choosing the students is not important, you must find the number of combinations of 20 students taken 4 at a time.

$$C(n, r) = \frac{n!}{(n-r)!r!} \quad \text{Combination formula}$$

$$C(20, 4) = \frac{20!}{(20-4)!4!} \quad n = 20, r = 4$$

$$= \frac{20!}{16!4!} \quad \text{or } 4845$$

There are 4845 possible ways to choose 4 students.

Example 2 **In how many ways can you choose 1 vowel and 2 consonants from a set of 26 letter tiles? (Assume there are 5 vowels and 21 consonants.)**

By the Fundamental Counting Principle, you can multiply the number of ways to select one vowel and the number of ways to select 2 consonants. Only the letters chosen matter, not the order in which they were chosen, so use combinations.

$C(5, 1)$ One of 5 vowels are drawn.

$C(21, 2)$ Two of 21 consonants are drawn.

$$C(5, 1) \cdot C(21, 2) = \frac{5!}{(5-1)!1!} \cdot \frac{21!}{(21-2)!2!} \quad \text{Combination formula}$$

$$= \frac{5!}{4!} \cdot \frac{21!}{19!2!} \quad \text{Simplify.}$$

$$= 5 \cdot 210 \text{ or } 1050$$

There are 1050 combinations of 1 vowel and 2 consonants.

Exercises

Evaluate each expression.

1. $C(5, 3)$ **10**
2. $C(7, 4)$ **35**
3. $C(15, 7)$ **6435**
4. $C(10, 5)$ **252**

5. **PLAYING CARDS** From a standard deck of 52 cards, in how many ways can 5 cards be drawn? **2,598,960**

6. **HOCKEY** How many hockey teams of 6 players can be formed from 14 players without regard to position played? **3003**

7. **COMMITTEES** From a group of 10 men and 12 women, how many committees of 5 men and 6 women can be formed? **232,848**

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____ DATE _____ PERIOD _____

12-2 Practice

Permutations and Combinations

Evaluate each expression.

- $P(8, 6)$ **20,160**
- $P(9, 7)$ **181,440**
- $P(3, 3)$ **6**
- $P(4, 3)$ **24**
- $P(4, 1)$ **4**
- $C(8, 2)$ **28**
- $C(11, 3)$ **165**
- $C(20, 18)$ **190**
- $C(9, 9)$ **1**
- $C(3, 1)$ **3**
- $C(9, 3) \cdot C(6, 2)$ **1260**

Determine whether each situation involves a *permutation* or a *combination*. Then find the number of possibilities.

- selecting a 4-person bobsled team from a group of 9 athletes
combination; 126
- an arrangement of the letters in the word *Canada*
permutation; 120
- arranging 4 charms on a bracelet that has a clasp, a front, and a back
permutation; 24
- selecting 3 desserts from 10 desserts that are displayed on a dessert cart in a restaurant
combination; 120
- an arrangement of the letters in the word *annually*
permutation; 5040
- forming a 2-person sales team from a group of 12 salespeople
combination; 66
- making 5-sided polygons by choosing any 5 of 11 points located on a circle to be the vertices
combination; 462
- seating 5 men and 5 women alternately in a row, beginning with a woman
permutation; 14,400
- STUDENT GROUPS** Farmington High is planning its academic festival. All math classes will send 2 representatives to compete in the math bowl. How many different groups of students can be chosen from a class of 16 students? **120**
- PHOTOGRAPHY** A photographer is taking pictures of a bride and groom and their 6 attendants. If she takes photographs of 3 people in a group, how many different groups can she photograph? **56**
- AIRLINES** An airline is hiring 5 flight attendants. If 8 people apply for the job, how many different groups of 5 attendants can the airline hire? **56**
- SUBSCRIPTIONS** A school librarian would like to buy subscriptions to 7 new magazines. Her budget, however, will allow her to buy only 4 new subscriptions. How many different groups of 4 magazines can she choose from the 7 magazines? **35**

Chapter 12

16

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-2 Word Problem Practice

Permutations and Combinations

- WAITING IN LINE** When the 12 students in Mr. Jaybird's class go to lunch, they form a single file line. Does forming a line involve a permutation or a combination of the students?
A permutation

- ART** Isabel needs to select three different colors of construction paper to make a flag for a school project. She can choose from a selection of 15 different colors. In how many ways can she pick her colors?
455

- SUDOKU** A popular game called "Sudoku" involves square arrays of numbers. In a game of Sudoku, every entry is an integer between 1 and 9, inclusive. No number appears twice in any row or column.

7	1	8	6	9	4	2	3	5
9	2	5	7	3	1	6	4	8
4	6	3	8	5	2	7	9	1
5	9	2	1	7	3	4	8	6
8	3	1	4	6	5	9	2	7
9	7	4	2	8	9	5	1	3
3	4	9	5	1	7	8	6	2
2	8	7	3	4	6	1	5	9
1	5	6	9	2	8	3	7	4

For a game of Sudoku, how many different possibilities are there for the first row of numbers?
362,880

- NAMES** Hannah is curious to know how many different 6 letter sequences she can make using each of the letters of her name exactly once. For example, "HANNAH," "AAHHNN," and "NAHNAH" are all possible sequences. How many total sequences are possible?
90

METEORITES For Exercises 5 and 6, use the following information.

- Over the course of several years, Kendra managed to collect 7 meteorites. Each one is unique.
- For a school science fair, Kendra displays her meteorites in a row. How many ways are there to order the meteorites?
5040

- She decides to trade three of her meteorites for a telescope after the fair. How many ways can she pick out 3 meteorites from her collection?
35

Lesson 12-2

17

Chapter 12

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-3 Lesson Reading Guide

Probability

Get Ready for the Lesson

Read the introduction to Lesson 12-3 in your textbook.

What is the probability that a person will *not* be struck by lightning in a given year?
749,999
750,000

Read the Lesson

- Indicate whether each of the following statements is *true* or *false*.
 - If an event can never occur, its probability is a negative number. **false**
 - If an event is certain to happen, its probability is 1. **true**
 - If an event can succeed in s ways and fail in f ways, then the probability of success is $\frac{s}{f}$. **false**
 - If an event can succeed in s ways and fail in f ways, then the odds against the event are $s:f$. **false**
 - A probability distribution is a function in which the domain is the sample space of an experiment. **true**
- A weather forecast says that the chance of rain tomorrow is 40%.
 - Write the probability that it will rain tomorrow as a fraction in lowest terms. **$\frac{2}{5}$**
 - Write the probability that it will not rain tomorrow as a fraction in lowest terms. **$\frac{3}{5}$**
 - What are the odds in favor of rain? **2:3**
 - What are the odds against rain? **3:2**
- Refer to the table in Example 4 on page 646 in your textbook.
 - What other sum has the same probability as a sum of 11? **3**
 - What are the odds of rolling a sum of 8? **5:31**
 - What are the odds against rolling a sum of 9? **8:1**

Remember What You Learned

- A good way to remember something is to explain it to someone else. Suppose that your friend Roberto is having trouble remembering the difference between probability and odds. What would you tell him to help him remember this easily?

Sample answer: Probability gives the ratio of successes to the total number of outcomes, while odds gives the ratio of successes to failures.

Chapter 12

20

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-3 Study Guide and Intervention

Probability


Probability and Odds In probability, a desired outcome is called a **success**; any other outcome is called a **failure**.

Probability of Success and Failure	If an event can succeed in s ways and fail in f ways, then the probabilities of success, $P(S)$, and of failure, $P(F)$, are as follows: $P(S) = \frac{s}{s+f}$ and $P(F) = \frac{f}{s+f}$.
Definition of Odds	If an event can succeed in s ways and fail in f ways, then the odds of success and of failure are as follows: Odds of success = $s:f$ Odds of failure = $f:s$

Example 1 When 3 coins are tossed, what is the probability that at least 2 are heads?

You can use a tree diagram to find the sample space.

First Coin **Second Coin** **Third Coin** **Possible Outcomes**
 Of the 8 possible outcomes, 4 have at least 2 heads. So the probability of tossing at least 2 heads is $\frac{4}{8}$ or $\frac{1}{2}$.



Example 2 What is the probability of picking 4 fiction books and 2 biographies from a best-seller list that consists of 12 fiction books and 6 biographies?

By the Fundamental Counting Principle, the number of successes is $C(12, 4) \cdot C(6, 2)$. The total number of selections, $s + f$, of 6 books is $C(18, 6)$.

$$P(4 \text{ fiction, } 2 \text{ biography}) = \frac{C(12, 4) \cdot C(6, 2)}{C(18, 6)} \text{ or about } 0.40$$

The probability of selecting 4 fiction books and 2 biographies is about 40%.

Exercises

Find the odds of an event occurring, given the probability of the event.

- $\frac{3}{7}$ **3:4**
- $\frac{4}{5}$ **4:1**
- $\frac{2}{13}$ **2:11**
- $\frac{1}{16}$ **1:14**

Find the probability of an event occurring, given the odds of the event.

- 10:1 $\frac{10}{11}$
- 2:5 $\frac{2}{7}$
- 4:9 $\frac{4}{13}$
- 8:3 $\frac{8}{11}$

One bag of candy contains 15 red candies, 10 yellow candies, and 6 green candies. Find the probability of each selection.

- picking a red candy $\frac{15}{31}$
- not picking a yellow candy $\frac{21}{31}$
- picking a green candy $\frac{6}{31}$
- not picking a red candy $\frac{16}{31}$

Chapter 12

20

Glencoe Algebra 2

Chapter 12

21

Glencoe Algebra 2

12-3 Study Guide and Intervention

(continued)

Probability

Probability Distributions A random variable is a variable whose value is the numerical outcome of a random event. A **probability distribution** for a particular random variable is a function that maps the sample space to the probabilities of the outcomes in the sample space.


Example

Suppose two dice are rolled. The table and the relative-frequency histogram show the distribution of the absolute value of the difference of the numbers rolled. Use the graph to determine which outcome is the most likely. What is its probability?

Difference	0	1	2	3	4	5
Probability	$\frac{1}{6}$	$\frac{5}{18}$	$\frac{2}{9}$	$\frac{1}{6}$	$\frac{1}{9}$	$\frac{1}{18}$

The greatest probability in the graph is $\frac{5}{18}$.

The most likely outcome is a difference of 1 and its probability is $\frac{5}{18}$.




Exercises

Four coins are tossed.

1. Complete the table below to show the probability distribution of the number of heads.

Number of Heads	0	1	2	3	4
Probability	$\frac{1}{16}$	$\frac{1}{4}$	$\frac{3}{8}$	$\frac{1}{4}$	$\frac{1}{16}$

2. Make relative-frequency distribution of the data.



12-3 Skills Practice

Probability

Ahmed is posting 2 photographs on his website. He has narrowed his choices to 4 landscape photographs and 3 portraits. If he chooses the two photographs at random, find the probability of each selection.

- $\frac{1}{7}$
- $\frac{2}{7}$
- $\frac{4}{7}$
- $\frac{20}{117}$
- $\frac{88}{273}$
- $\frac{40}{91}$
- $\frac{20}{117}$
- $\frac{88}{273}$
- $\frac{2}{7}$
- $\frac{4}{7}$

The Carubas have a collection of 28 video movies, including 12 westerns and 16 science fiction. Elise selects 3 of the movies at random to bring to a sleep-over at her friend's house. Find the probability of each selection.

- $\frac{55}{819}$
- $\frac{20}{117}$
- $\frac{88}{273}$
- $\frac{40}{91}$
- $\frac{20}{117}$
- $\frac{40}{91}$
- $\frac{20}{117}$
- $\frac{88}{273}$
- $\frac{2}{7}$
- $\frac{4}{7}$

For Exercises 10–13, use the chart that shows the class and gender statistics for the students taking an Algebra 1 or Algebra 2 class at La Mesa High School.

If a student taking Algebra 1 or Algebra 2 is selected at random, find each probability. Express as decimals rounded to the nearest thousandth.

- $P(\text{sophomore/female})$ **0.208**
- $P(\text{junior/male})$ **0.143**
- $P(\text{freshman/male})$ **0.136**
- $P(\text{freshman/female})$ **0.145**

Class/Gender	Number
Freshman/Male	95
Freshman/Female	101
Sophomore/Male	154
Sophomore/Female	145
Junior/Male	100
Junior/Female	102

Find the odds of an event occurring, given the probability of the event.

- $\frac{5}{8}$ **5:3**
- $\frac{2}{7}$ **2:5**
- $\frac{3}{9}$ **3:2**
- $\frac{1}{10}$ **1:9**
- $\frac{5}{6}$ **5:1**
- $\frac{5}{12}$ **5:7**

Find the probability of an event occurring, given the odds of the event.

- 2:1 $\frac{2}{3}$
- 8:9 $\frac{8}{17}$
- 4:1 $\frac{4}{5}$
- 1:9 $\frac{1}{10}$
- 2:7 $\frac{2}{9}$
- 5:9 $\frac{5}{14}$

NAME _____ DATE _____ PERIOD _____

12-3 Practice

Probability

A bag contains 1 green, 4 red, and 5 yellow balls. Two balls are selected at random. Find the probability of each selection.

1. $P(2 \text{ red})$ $\frac{2}{15}$
2. $P(1 \text{ red and 1 yellow})$ $\frac{4}{9}$
3. $P(1 \text{ green and 1 yellow})$ $\frac{1}{9}$
4. $P(2 \text{ green})$ 0
5. $P(2 \text{ red and 1 yellow})$ 0
6. $P(1 \text{ red and 1 green})$ $\frac{4}{45}$

A bank contains 3 pennies, 8 nickels, 4 dimes, and 10 quarters. Two coins are selected at random. Find the probability of each selection.

7. $P(2 \text{ pennies})$ $\frac{1}{100}$
8. $P(2 \text{ dimes})$ $\frac{1}{50}$
9. $P(1 \text{ nickel and 1 dime})$ $\frac{8}{75}$
10. $P(1 \text{ quarter and 1 penny})$ $\frac{4}{15}$
11. $P(1 \text{ quarter and 1 nickel})$ 0
12. $P(2 \text{ dimes and 1 quarter})$ 0

Henrico visits a home decorating store to choose wallpaper for his new house. The store has 28 books of wallpaper samples, including 10 books of WallPride samples and 18 books of Deluxe Wall Coverings samples. The store will allow Henrico to bring 4 books home for a few days so he can decide which wallpapers he wants to buy. If Henrico randomly chooses 4 books to bring home, find the probability of each selection.

13. $P(4 \text{ WallPride})$ $\frac{2}{195}$
14. $P(2 \text{ WallPride and 2 Deluxe})$ $\frac{153}{455}$
15. $P(1 \text{ WallPride and 3 Deluxe})$ $\frac{544}{1365}$
16. $P(3 \text{ WallPride and 1 Deluxe})$ $\frac{48}{455}$

For Exercises 17–20, use the table that shows the range of verbal SAT scores for freshmen at a small liberal arts college. If a freshman student is chosen at random, find each probability. Express as decimals rounded to the nearest thousandth.

Range	400–449	450–499	500–549	550–559	600–649	650+
Number of Students	129	275	438	602	620	412

17. $P(400-449)$ 0.052
18. $P(550-559)$ 0.243
19. $P(\text{at least } 650)$ 0.166

Find the odds of an event occurring, given the probability of the event.

20. $\frac{4}{4}$ $4:7$
21. $\frac{12}{13}$ $12:1$
22. $\frac{5}{99}$ $5:94$
23. $\frac{1}{1000}$
24. $\frac{5}{16}$ $5:11$
25. $\frac{3}{95}$ $3:92$
26. $\frac{9}{70}$ $9:61$
27. $\frac{8}{15}$ $8:7$

Find the probability of an event occurring, given the odds of the event.

28. 2:23 $\frac{2}{25}$
29. 2:5 $\frac{2}{7}$
30. 15:1 $\frac{15}{16}$
31. 9:7 $\frac{9}{16}$
32. 11:14 $\frac{11}{25}$
33. 1000:1 $\frac{1000}{1001}$
34. 12:17 $\frac{12}{29}$
35. 8:13 $\frac{8}{21}$

Chapter 12

24

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-3 Word Problem Practice

Probability

1. **ART** The letters “A”, “R”, and “T” are written on three different pieces of paper. The pieces of paper are then put in a bag and mixed up. Logan picks each letter without looking and places them side by side. What is the probability that the letters spell “ART”?

$\frac{1}{6}$

2. **AGE** There are 24 students in Miss Mason’s third grade class, all born on different days. Eleven students are boys. In the morning, the classroom is empty. One student arrives followed by another. What is the probability that when the first two students arrive, one is a boy and the other a girl?

52%

3. **DICE** Jamal rolls two six-sided dice, one after the other. What is the probability that the second die shows a number larger than the first die?

$\frac{5}{12}$

4. **LANGUAGES** Noah cannot decide whether to learn French, German, Italian, Russian, or Chinese. He assigns each language a different number from 0 to 4. He then takes four fair coins and flips them. He decided to take the language corresponding to the number of coins that come up heads. Does Noah’s method for choosing a language give each language the same chance of being chosen? Explain.

No. For example, the probability of getting 0 is 1 in 16 but the probability of getting 1 is 1 in 4.

ICE CREAM For Exercises 5–7, use the following information.

A survey of the students in Mr. Orr’s fifth grade class asked each student to name their favorite flavor of ice cream. The results are shown in the table below.

Flavor	Number of Students
Vanilla	10
Chocolate	9
Butternut	5
Strawberry	4
Banana	1
Coffee	1

5. A student from Mr. Orr’s class is selected at random. What is the probability that the student’s favorite flavor of ice cream is chocolate?

$\frac{3}{10}$

6. A student from Mr. Orr’s class is selected at random. What is the probability that the student’s favorite flavor of ice cream is banana?

$\frac{1}{30}$

7. A student from Mr. Orr’s class is selected at random. Is it more likely that the student prefers either butternut or strawberry or that the student prefers either chocolate or banana?

Chocolate or banana is more likely.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Chapter 12

25

Glencoe Algebra 2

NAME _____

DATE _____

PERIOD _____

12-3 Enrichment

Geometric Probability

If a dart, thrown at random, hits the triangular board shown at the right, what is the chance that it will hit the shaded region? This chance, also called a probability, can be determined by comparing the area of the shaded region to the area of the board. This ratio indicates what fraction of the tosses should hit in the shaded region.


$$\frac{\text{area of shaded region}}{\text{area of triangular board}} = \frac{\frac{1}{2}(4)(6)}{\frac{1}{2}(8)(6)}$$

$$= \frac{12}{24} \text{ or } \frac{1}{2}$$

In general, if S is a subregion of some region R , then the probability, $P(S)$, that a point, chosen at random, belongs to subregion S is given by the following.

$$P(S) = \frac{\text{area of subregion } S}{\text{area of region } R}$$


Find the probability that a point, chosen at random, belongs to the shaded subregions of the following regions.



1

$$\frac{1}{2}$$


2.



3.

$$\frac{5}{9}$$

4.



5.

$$\frac{\pi}{4}$$

The dart board shown at the right has 5 concentric circles whose centers are also the center of the square board. Each side of the board is 38 cm, and the radii of the circles are 2 cm, 5 cm, 8 cm, 11 cm, and 14 cm. A dart hitting within one of the circular regions scores the number of points indicated on the board, while a hit anywhere else scores 0 points. If a dart, thrown at random, hits the board, find the probability of scoring the indicated number of points.

4. 0 points
 $\frac{361 - 49\pi}{361}$


5. 1 point
 $\frac{75\pi}{1444}$

7. 3 points
 $\frac{39\pi}{1444}$

8. 4 points
 $\frac{21\pi}{1444}$

6. 2 points
 $\frac{57\pi}{1444}$

9. 5 points
 $\frac{\pi}{361}$



NAME _____

DATE _____

PERIOD _____

12-4 Lesson Reading Guide

Multiplying Probabilities

Get Ready for the Lesson

Read the introduction to Lesson 12-4 in your textbook.

Write the probability that Yao Ming made a field goal shot during the 2004–05 season as a fraction in lowest terms. (Your answer should not include a decimal.) $\frac{276}{5}$

Read the Lesson

1. A bag contains 4 yellow balls, 5 red balls, 1 white ball, and 2 black balls. A ball is drawn from the bag and is not replaced. A second ball is drawn.

- Let Y be the event “first ball is yellow” and B be the event “second ball is black.” Are these events *independent* or *dependent*? **dependent**
- Tell which formula you would use to find the probability that the first ball is yellow and the second ball is black. **C**

A. $P(Y \text{ and } B) = \frac{P(Y)}{P(Y) + P(B)}$

B. $P(Y \text{ and } B) = P(Y) \cdot P(B)$

C. $P(Y \text{ and } B) = P(Y) \cdot P(B \text{ following } Y)$

c. Which equation shows the correct calculation of this probability? **B**

A. $\frac{1}{3} + \frac{2}{11} = \frac{17}{33}$

B. $\frac{1}{3} \cdot \frac{2}{11} = \frac{2}{33}$

C. $\frac{1}{3} + \frac{1}{6} = \frac{1}{2}$

D. $\frac{1}{3} \cdot \frac{1}{6} = \frac{1}{18}$

d. Which equation shows the correct calculation of the probability that if three balls are drawn in succession without replacement, all three will be red? **B**

A. $\frac{5}{12} \cdot \frac{5}{12} \cdot \frac{5}{12} = \frac{125}{1728}$

B. $\frac{5}{12} \cdot \frac{4}{11} \cdot \frac{3}{10} = \frac{22}{660}$

C. $\frac{5}{12} + \frac{4}{11} + \frac{3}{10} = \frac{713}{660}$

Remember What You Learned

2. Some students have trouble remembering a lot of formulas, so they try to keep the number of formulas they have to know to a minimum. Can you learn just one formula that will allow you to find probabilities for both independent and dependent events? Explain your reasoning. **Sample answer: Just remember the formula for dependent events: $P(A \text{ and } B) = P(A) \cdot P(B \text{ following } A)$. When the events are independent, $P(A \text{ and } B) = P(A) \cdot P(B)$, so the formula for dependent events simplifies to $P(A \text{ and } B) = P(A) \cdot P(B)$, which is the correct formula for independent events.**

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____

DATE _____

PERIOD _____


12-4 Study Guide and Intervention

Multiplying Probabilities

Probability of Independent Events

Probability of Two Independent Events
 If two events, A and B , are independent, then the probability of both occurring is $P(A \text{ and } B) = P(A) \cdot P(B)$.

Example 1 In a board game, each player has 3 different-colored markers. To move around the board, the player first spins a spinner to determine which piece can be moved. He or she then rolls a die to determine how many spaces that colored piece should move. On a given turn, what is the probability that a player will be able to move the yellow piece more than 2 spaces?



Let A be the event that the spinner lands on yellow, and let B be the event that the die shows a number greater than 2. The probability of A is $\frac{1}{3}$, and the probability of B is $\frac{2}{3}$.

$$P(A \text{ and } B) = P(A) \cdot P(B) \quad \text{Probability of independent events}$$

$$= \frac{1}{3} \cdot \frac{2}{3} \text{ or } \frac{2}{9} \quad \text{Substitute and multiply.}$$

The probability that the player can move the yellow piece more than 2 spaces is $\frac{2}{9}$.

Exercises

A die is rolled 3 times. Find the probability of each event.

- a 1 is rolled, then a 2, then a 3 $\frac{1}{216}$
- a 1 or a 2 is rolled, then a 3, then a 5 or a 6 $\frac{1}{54}$
- 2 odd numbers are rolled, then a 6 $\frac{1}{24}$
- a number less than 3 is rolled, then a 3, then a number greater than 3 $\frac{1}{36}$
- A box contains 5 triangles, 6 circles, and 4 squares. If a figure is removed, replaced, and a second figure is picked, what is the probability that a triangle and then a circle will be picked? $\frac{2}{15}$ or about 0.13
- A bag contains 5 red marbles and 4 white marbles. A marble is selected from the bag, then replaced, and a second selection is made. What is the probability of selecting 2 red marbles? $\frac{25}{81}$ or about 0.31
- A jar contains 7 lemon jawbreakers, 3 cherry jawbreakers, and 8 rainbow jawbreakers. What is the probability of selecting 2 lemon jawbreakers in succession providing the jawbreaker drawn first is then replaced before the second is drawn?
 $\frac{49}{324}$ or about 0.15

Chapter 12

28

Glencoe Algebra 2

NAME _____

DATE _____

PERIOD _____

12-4 Study Guide and Intervention

Multiplying Probabilities

Probability of Dependent Events

Probability of Two Dependent Events
 If two events, A and B , are dependent, then the probability of both events occurring is $P(A \text{ and } B) = P(A) \cdot P(B \text{ following } A)$.

Example 1 There are 7 dimes and 9 pennies in a wallet. Suppose two coins are to be selected at random, without replacing the first one. Find the probability of picking a penny and then a dime.

Because the coin is not replaced, the events are dependent.

Thus, $P(A \text{ and } B) = P(A) \cdot P(B \text{ following } A)$.

$P(\text{penny, then dime}) = P(\text{penny}) \cdot P(\text{dime following penny})$

$$\frac{9}{16} \cdot \frac{7}{15} = \frac{21}{80}$$

The probability is $\frac{21}{80}$ or about 0.26

Example 2 What is the probability of drawing, without replacement, 3 hearts, then a spade from a standard deck of cards?

Since the cards are not replaced, the events are dependent. Let H represent a heart and S represent a spade.

$$P(H, H, H, S) = P(H) \cdot P(H \text{ following } H) \cdot P(H \text{ following } 2 Hs) \cdot P(S \text{ following } 3 Hs)$$

$$= \frac{13}{52} \cdot \frac{12}{51} \cdot \frac{11}{50} \cdot \frac{13}{49} \text{ or about } 0.003$$

The probability is about 0.003 of drawing 3 hearts, then a spade.

Exercises

Find each probability.

- The cup on Sophie's desk holds 4 red pens and 7 black pens. What is the probability of her selecting first a black pen, then a red one? $\frac{14}{55}$ or about 0.25
- What is the probability of drawing two cards showing odd numbers from a set of cards that show the first 20 counting numbers if the first card is not replaced before the second is chosen? $\frac{9}{38}$ or about 0.24
- There are 3 quarters, 4 dimes, and 7 nickels in a change purse. Suppose 3 coins are selected without replacement. What is the probability of selecting a quarter, then a dime, and then a nickel? $\frac{1}{26}$ or about 0.04
- A basket contains 4 plums, 6 peaches, and 5 oranges. What is the probability of picking 2 oranges, then a peach if 3 pieces of fruit are selected at random? $\frac{4}{91}$ or about 0.04
- A photographer has taken 8 black and white photographs and 10 color photographs for a brochure. If 4 photographs are selected at random, what is the probability of picking first 2 black and white photographs, then 2 color photographs? $\frac{7}{102}$ or about 0.07

Chapter 12

29

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-4 Practice

Multiplying Probabilities

A die is rolled three times. Find each probability.

- $P(\text{three 4s}) = \frac{1}{216}$
- $P(\text{no 4s}) = \frac{125}{216}$
- $P(2, \text{ then } 3, \text{ then } 1) = \frac{1}{216}$
- $P(\text{three different even numbers}) = \frac{1}{36}$
- $P(\text{any number, then 5, then 5}) = \frac{1}{36}$
- $P(\text{even number, then odd number, then } 1) = \frac{1}{24}$

There are 3 nickels, 2 dimes, and 5 quarters in a purse. Three coins are selected in succession at random. Find the probability.

- $P(\text{nickel, then dime, then quarter})$, if no replacement occurs $\frac{3}{100}$
- $P(\text{nickel, then dime, then quarter})$, if replacement occurs $\frac{1}{24}$
- $P(2 \text{ nickels, then } 1 \text{ quarter})$, if no replacement occurs $\frac{1}{24}$
- $P(2 \text{ nickels, then } 1 \text{ quarter})$, if replacement occurs $\frac{1}{24}$
- $P(3 \text{ dimes})$, if replacement occurs $\frac{1}{125}$
- $P(3 \text{ dimes})$, if no replacement occurs 0

For Exercises 12 and 13, determine whether the events are independent or dependent. Then find each probability.


- Serena is creating a painting. She wants to use 2 more colors. She chooses randomly from 6 shades of red, 10 shades of green, 4 shades of yellow, 4 shades of purple, and 6 shades of blue. What is the probability that she chooses 2 shades of green? **dependent; $\frac{3}{29}$**
- Kershels mother is shopping at a bakery. The owner offers Kershel a cookie from a jar containing 22 chocolate chip cookies, 18 sugar cookies, and 15 oatmeal cookies. Without looking, Kershel selects one, drops it back in, and then randomly selects another. What is the probability that neither selection was a chocolate chip cookie? **independent; $\frac{9}{9}$**
- METEOROLOGY** The Fadeeva's are planning a 3-day vacation to the mountains. A long-range forecast reports that the probability of rain each day is 10%. Assuming that the daily probabilities of rain are independent, what is the probability that there is no rain on the first two days, but that it rains on the third day? **$\frac{81}{1000}$**

RANDOM NUMBERS For Exercises 15 and 16, use the following information.

Anita has a list of 20 jobs around the house to do, and plans to do 3 of them today. She assigns each job a number from 1 to 20, and sets her calculator to generate random numbers from 1 to 20, which can reoccur. Of the jobs, 3 are outside, and the rest are inside.

15. Sketch a tree diagram showing all of the possibilities that the first three numbers generated correspond to inside jobs or outside jobs. Use it to find the probability that the first two numbers correspond to inside jobs, and the third to an outside job. **0.106375**

16. What is the probability that the number generated corresponds to an outside job three times in a row? **0.003375**



NAME _____ DATE _____ PERIOD _____

12-4 Skills Practice

Multiplying Probabilities

A die is rolled twice. Find each probability.

- $P(5, \text{ then } 6) = \frac{1}{36}$
- $P(\text{no 2s}) = \frac{25}{36}$
- $P(\text{two 1s}) = \frac{1}{36}$
- $P(\text{not } 1, \text{ then not } 2) = \frac{25}{36}$
- $P(4, \text{ then not } 6) = \frac{5}{36}$

A board game uses a set of 6 different cards. Each card displays one of the following figures: a star, a square, a circle, a diamond, a rectangle, or a pentagon. The cards are placed face down, and a player chooses two cards. Find each probability.

- $P(\text{circle, then star})$, if no replacement occurs $\frac{1}{30}$
- $P(\text{diamond, then square})$, if replacement occurs $\frac{1}{36}$
- $P(2 \text{ polygons})$, if replacement occurs $\frac{25}{36}$
- $P(2 \text{ polygons})$, if no replacement occurs $\frac{2}{3}$
- $P(\text{circle, then hexagon})$, if no replacement occurs 0

Determine whether the events are independent or dependent. Then find each probability.



- A mixed box of herbal teabags contains 2 lemon teabags, 3 orange-mango teabags, 3 chamomile teabags, and 1 apricot-ginger teabag. Kevin chooses 2 teabags at random to bring to work with him. What is the probability that he first chooses a lemon teabag and then a chamomile teabag? **dependent; $\frac{1}{12}$**
- The chart shows the selection of olive oils that Hasha finds in a specialty foods catalog. If she randomly selects one type of oil, then randomly selects another, different oil, what is the probability that both selections are domestic, first cold pressed oils? **dependent; $\frac{21}{820}$**

Type of Oil	Domestic	Imported
Pure	2	5
Cold Pressed	4	8
First Cold Pressed	7	15

For Exercises 14 and 15, two thirds of the area of the spinner earns you 50 points. Suppose you spin the spinner twice.

14. Sketch a tree diagram showing all of the possibilities. Use it to find the probability of spinning 50 points, then 100 points. **$\frac{2}{9}$**

15. What is the probability that you get 100 points on each spin? **$\frac{1}{9}$**



NAME _____ DATE _____ PERIOD _____

12-4 Word Problem Practice

Multiplying Probabilities

- BUSSING** Portia and Quinton use the same bus stop when they go to work. They arrive at the bus stop independently of each other. The probability that Portia catches the 7:45 A.M. bus is $\frac{3}{5}$. The probability that Quinton catches the 7:45 A.M. bus is $\frac{1}{2}$. What is the probability that they both catch the 7:45 A.M. bus on the same day?



What is the probability that she does not pick the "win" card on her first try but does find it with her second?

$\frac{1}{4}$

WALLETS For Exercises 5 and 6, use the following information.

- Wayne has 1 ten-dollar bill, 2 five-dollar bills, and 5 one-dollar bills in his wallet.
5. Wayne randomly chooses a bill from his wallet, puts it back, then picks another bill, and puts that one back, too. What is the probability that both were five-dollar bills?
- $\frac{1}{16}$
6. Wayne randomly pulls out a bill from his wallet, and then, without putting it back, randomly pulls a second bill from his wallet. He then puts both bills back into the wallet. What is the probability that both of the bills pulled out were five-dollar bills?
- $\frac{1}{28}$

- GOODY BAGS** Ryan and Sophia are given goody bags with identical contents. The probability of reaching into either of these goody bags and pulling out a stick of chewing gum is $\frac{1}{10}$. Ryan and Sophia each reach into their own goody bag and randomly pull out something. What is the probability that they both pulled out a stick of chewing gum?
- $\frac{1}{100}$
- PENCILS** A box of pencils contains 11 type 2 pencils and 5 type 3 pencils. Tara picks out a pencil from the box without looking and keeps it. Then, Upton picks out a pencil from the box without looking. What is the probability that Tara picks a type 2 pencil and Upton picks a type 3 pencil?
- $\frac{11}{48}$

NAME _____ DATE _____ PERIOD _____

12-4 Enrichment

Conditional Probability

Suppose a pair of dice is thrown. It is known that the sum is greater than seven. Find the probability that the dice match.

The probability of an event given the occurrence of another event is called *conditional probability*. The conditional probability of event A , the dice match, given event B , their sum is greater than seven, is denoted $P(A|B)$.

There are 15 sums greater than seven and there are 36 possible pairs altogether.

$P(B) = \frac{15}{36}$

There are three matching pairs greater than seven.

$P(A \text{ and } B) = \frac{3}{36}$

$$P(A|B) = \frac{P(A \text{ and } B)}{P(B)}$$

$$P(A|B) = \frac{\frac{3}{36}}{\frac{15}{36}} \text{ or } \frac{1}{5}$$

The conditional probability is $\frac{1}{5}$.

A card is drawn from a standard deck of 52 and is found to be red. Given that event, find each of the following probabilities.

- $P(\text{heart})$ $\frac{1}{2}$
- $P(\text{ace})$ $\frac{1}{13}$
- $P(\text{face card})$ $\frac{3}{13}$
- $P(\text{jack or ten})$ $\frac{2}{13}$
- $P(\text{six of spades})$ 0
- $P(\text{six of hearts})$ $\frac{1}{26}$

A sports survey taken at Stirers High School shows that 48% of the respondents liked soccer, 66% liked basketball, and 38% liked hockey. Also, 30% liked soccer and basketball, 22% liked basketball and hockey and 28% liked soccer and hockey. Finally, 12% liked all three sports. Find each of the following probabilities.

- The probability Meg likes soccer if she likes basketball. $\frac{30}{66}$ or $\frac{5}{11}$
- The probability Biff likes basketball if he likes soccer. $\frac{30}{48}$ or $\frac{5}{8}$
- The probability Muffy likes hockey if she likes basketball. $\frac{22}{66}$ or $\frac{1}{3}$
- The probability Greg likes soccer and basketball if he likes soccer. $\frac{12}{48}$ or $\frac{1}{4}$

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____ DATE _____ PERIOD _____

12-5 Study Guide and Intervention

Adding Probabilities

Mutually Exclusive Events Events that cannot occur at the same time are called mutually exclusive events.

Probability of Mutually Exclusive Events
 $P(A \text{ or } B) = P(A) + P(B)$.
 If two events, A and B, are mutually exclusive, then

This formula can be extended to any number of mutually exclusive events.

Example 1 To choose an afternoon activity, summer campers pull slips of paper out of a hat. Today there are 25 slips for a nature walk, 35 slips for swimming, and 30 slips for arts and crafts. What is the probability that a camper will pull a slip for a nature walk or for swimming?

These are mutually exclusive events. Note that there is a total of 90 slips.

$$P(\text{nature walk or swimming}) = P(\text{nature walk}) + P(\text{swimming})$$

$$= \frac{25}{90} + \frac{35}{90} \text{ or } \frac{2}{3}$$

The probability of a camper's pulling out a slip for a nature walk or for swimming is $\frac{2}{3}$.

Example 2 By the time one tent of 6 campers gets to the front of the line, there are only 10 nature walk slips and 15 swimming slips left. What is the probability that more than 4 of the 6 campers will choose a swimming slip?

$$P(\text{more than 4 swimmers}) = P(5 \text{ swimmers}) + P(6 \text{ swimmers})$$

$$= \frac{C(10, 1) \cdot C(15, 5)}{C(25, 6)} + \frac{C(10, 0) \cdot C(15, 6)}{C(25, 6)}$$

$$\approx 0.2$$

The probability of more than 4 of the campers swimming is about 0.2.

Exercises

Find each probability.

- A bag contains 45 dyed eggs: 15 yellow, 12 green, and 18 red. What is the probability of selecting a green or a red egg? $\frac{2}{3}$
- The letters from the words LOVE and LIVE are placed on cards and put in a box. What is the probability of selecting an L or an O from the box? $\frac{3}{8}$
- A pair of dice is rolled, and the two numbers are added. What is the probability that the sum is either a 5 or a 7? $\frac{5}{18}$ or about 0.28
- A bowl has 10 whole wheat crackers, 16 sesame crackers, and 14 rye crisps. If a person picks a cracker at random, what is the probability of picking either a sesame cracker or a rye crisp? $\frac{3}{4}$
- An art box contains 12 colored pencils and 20 pastels. If 5 drawing implements are chosen at random, what is the probability that at least 4 of them are pastels? about 0.37

Chapter 12

35

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-5 Lesson Reading Guide

Adding Probabilities

Get Ready for the Lesson

Read the introduction to Lesson 12-5 in your textbook.

Why do the percentages shown on the bar graph add up to more than 100%? **Sample answer: Many teens do one or more of the listed online activities.**

Read the Lesson

1. Indicate whether the events in each pair are *inclusive* or *mutually exclusive*.

- drawing a queen from a standard deck of cards
 drawing a diamond from a standard deck of cards **inclusive**
- drawing a jack from a standard deck of cards
 drawing a king from a standard deck of cards **mutually exclusive**

2. Marla took a quiz on this lesson that contained the following problem.

Each of the integers from 1 through 25 is written on a slip of paper and placed in an envelope. If one slip is drawn at random, what is the probability that it is odd or a multiple of 5?

Here is Marla's work.

$$P(\text{odd}) = \frac{13}{25} \quad P(\text{multiple of 5}) = \frac{5}{25} \text{ or } \frac{1}{5}$$

$$P(\text{odd or multiple of 5}) = P(\text{odd}) + P(\text{multiple of 5})$$

$$= \frac{13}{25} + \frac{5}{25} = \frac{18}{25}$$

a. Why is Marla's work incorrect? **Sample answer: Marla used the formula for mutually exclusive events, but the events are inclusive. She should use the formula for inclusive events so that the odd multiples of 5 will not be counted twice.**

b. Show the corrected work.

$$P(\text{odd or multiple of 5}) = P(\text{odd}) + P(\text{multiple of 5}) - P(\text{odd multiple of 5})$$

$$= \frac{13}{25} + \frac{5}{25} - \frac{3}{25} = \frac{15}{25} = \frac{3}{5}$$

Remember What You Learned

- Some students have trouble remembering a lot of formulas, so they try to keep the number of formulas they have to know to a minimum. Can you learn just one formula that will allow you to find probabilities for both mutually exclusive and inclusive events? Explain your reasoning. **Sample answer: Just remember the formula for inclusive events: $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$. When the events are mutually exclusive, $P(A \text{ and } B) = 0$, so the formula for inclusive events simplifies to $P(A \text{ and } B) = P(A) + P(B)$, which is the correct formula for mutually exclusive events.**

Chapter 12

34

Glencoe Algebra 2

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____ DATE _____ PERIOD _____

12-5 Study Guide and Intervention *(continued)*

Adding Probabilities

Inclusive Events

Probability of Inclusive Events If two events, A and B , are inclusive, $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$.

Example What is the probability of drawing a face card or a black card from a standard deck of cards?


The two events are inclusive, since a card can be both a face card and a black card.
 $P(\text{face card or black card}) = P(\text{face card}) + P(\text{black card}) - P(\text{black face card})$
 $= \frac{3}{13} + \frac{1}{2} - \frac{3}{26}$
 $= \frac{8}{13}$ or about 0.62

The probability of drawing either a face card or a black card is about 0.62

EXERCISES

Find each probability.

1. What is the probability of drawing a red card or an ace from a standard deck of cards?
 $\frac{7}{13}$ or about 0.54
2. Three cards are selected from a standard deck of 52 cards. What is the probability of selecting a king, a queen, or a red card?
 $\frac{15}{26}$ or about 0.58
3. The letters of the alphabet are placed in a bag. What is the probability of selecting a vowel or one of the letters from the word QUIZ?
 $\frac{7}{26}$ or about 0.27
4. A pair of dice is rolled. What is the probability that the sum is odd or a multiple of 3?
 $\frac{2}{3}$ or about 0.67



5. The Venn diagram at the right shows the number of juniors on varsity sports teams at Elmwood High School. Some athletes are on varsity teams for one season only, some athletes for two seasons, and some for all three seasons. If a varsity athlete is chosen at random from the junior class, what is the probability that he or she plays a fall or winter sport?
 $\frac{13}{16}$

Chapter 12

36

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-5 Skills Practice

Adding Probabilities

Eli has 10 baseball cards of 10 different players in his pocket. Three players are pitchers, 5 are outfielders, and 2 are catchers. If Eli randomly selects a card to trade, find each probability.

1. $P(\text{pitcher or outfielder})$ $\frac{4}{5}$
2. $P(\text{pitcher or catcher})$ $\frac{1}{2}$
3. $P(\text{outfielder or catcher})$ $\frac{7}{10}$

A die is rolled. Find each probability.

4. $P(5 \text{ or } 6)$ $\frac{1}{3}$
5. $P(\text{at least a } 3)$ $\frac{2}{3}$
6. $P(\text{less than } 4)$ $\frac{1}{2}$

Determine whether the events are *mutually exclusive* or *inclusive*. Then find the probability.

7. A die is rolled. What is the probability of rolling a 3 or a 4? **mutually exclusive; $\frac{1}{3}$**
8. A die is rolled. What is the probability of rolling an even number or a 4? **inclusive; $\frac{1}{2}$**
9. A card is drawn from a standard deck of cards. What is the probability of drawing a king or a queen? **mutually exclusive; $\frac{2}{13}$**
10. A card is drawn from a standard deck of cards. What is the probability of drawing a jack or a heart? **inclusive; $\frac{4}{13}$**
11. The sophomore class is selling Mother's Day plants to raise money. Susan's prize for being the top seller of plants is a choice of a book, a CD, or a video. She can choose from 6 books, 3 CDs, and 5 videos. What is the probability that Susan selects a book or a CD? **mutually exclusive; $\frac{9}{14}$**


A spinner numbered 1–10 is spun. Find each probability.

12. $P(\text{less than } 5 \text{ or even})$ $\frac{7}{10}$
13. $P(\text{even or odd})$ 1
14. $P(\text{prime or even})$ $\frac{4}{5}$

Two cards are drawn from a standard deck of cards. Find each probability.

15. $P(\text{both red or both black})$ $\frac{25}{51}$
16. $P(\text{both aces or both red})$ $\frac{55}{221}$
17. $P(\text{both } 2\text{s or both less than } 5)$ $\frac{11}{221}$
18. $P(\text{both black or both less than } 5)$ $\frac{188}{663}$

For Exercises 19 and 20, use the Venn diagram that shows the number of participants in two different kinds of aerobic exercise classes that are offered at a health club. Determine each probability if a person is selected at random from the participants.



19. $P(\text{step aerobics or jazzercise, but not both})$ $\frac{49}{62}$
20. $P(\text{step aerobics and jazzercise})$ $\frac{13}{62}$

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Chapter 12

37

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-5 Word Problem Practice

Adding Probabilities

- 1. PICK-UP** When Tina's parents pick her up from school, there is a $\frac{1}{5}$ chance that she will be in the library, a $\frac{1}{2}$ chance that she will be on the playground, and a $\frac{3}{10}$ chance that she will be in her classroom. What is the probability that when Tina's parents pick her up, she is found in her classroom or on the playground?
- 4** **5** **3**

PASSENGERS For Exercises 5 and 6, use the following information.

On an airplane flight, some passengers travel with carry-on luggage while others travel with a suitcase. Some passengers travel with carry-on luggage and a suitcase. Everyone travels with some form of luggage.

- 5.** On one flight, there was no passenger with both carry-on luggage and a suitcase. On this flight are the events of picking a passenger with carry-on luggage and picking a passenger with a suitcase mutually exclusive?
- Yes**

- 6.** On another flight, there are 120 passengers. Of those 120 passengers, 80 have carry-on luggage and 70 have a suitcase. What is the probability that a passenger has both carry-on luggage and a suitcase?
- $\frac{1}{4}$**

NAME _____ DATE _____ PERIOD _____


12-5 Practice

Adding Probabilities

- An urn contains 7 white marbles and 5 blue marbles. Four marbles are selected without replacement. Find each probability.
- 1.** $P(4 \text{ white or } 4 \text{ blue})$ **$\frac{8}{99}$** **2.** $P(\text{exactly } 3 \text{ white})$ **$\frac{35}{99}$** **3.** $P(\text{at least } 3 \text{ white})$ **$\frac{14}{33}$**
4. $P(\text{fewer than } 3 \text{ white})$ **$\frac{19}{33}$** **5.** $P(3 \text{ white or } 3 \text{ blue})$ **$\frac{49}{99}$** **6.** $P(\text{no white or no blue})$ **$\frac{8}{99}$**
- Jason and Maria are playing a board game in which three dice are tossed to determine a player's move. Find each probability.**
- 7.** $P(\text{no } 5\text{s})$ **$\frac{5}{72}$** **8.** $P(\text{three } 5\text{s})$ **$\frac{1}{216}$** **9.** $P(\text{at least two } 5\text{s})$ **$\frac{2}{27}$**
10. $P(\text{no } 5\text{s})$ **$\frac{125}{216}$** **11.** $P(\text{one } 5)$ **$\frac{25}{72}$** **12.** $P(\text{one } 5 \text{ or two } 5\text{s})$ **$\frac{5}{12}$**

Determine whether the events are mutually exclusive or inclusive. Then find the probability.

- 13.** A clerk chooses 4 CD players at random for floor displays from a shipment of 24 CD players. If 15 of the players have a blue case and the rest have a red case, what is the probability of choosing 4 players with a blue case or 4 players with a red case? **mutual. exclus.; $\frac{71}{72}$**
- 14.** A department store employs 28 high school students, all juniors and seniors. Six of the 12 seniors are females and 12 of the juniors are males. One student employee is chosen at random. What is the probability of selecting a senior or a female? **inclusive; $\frac{4}{7}$**
- 15.** A restaurant has 5 pieces of apple pie, 4 pieces of chocolate cream pie, and 3 pieces of blueberry pie. If Janine selects a piece of pie at random for dessert, what is the probability that she selects either apple or chocolate cream? **mutually exclusive; $\frac{3}{7}$**
- 16.** At a statewide meeting, there are 20 school superintendents, 13 principals, and 6 assistant principals. If one of these people is chosen at random, what is the probability that he or she is either a principal or an assistant principal? **mutually exclusive; $\frac{19}{39}$**
- 17.** An airline has one bank of 13 telephones at a reservations office. Of the 13 operators who work there, 8 take reservations for domestic flights and 5 take reservations for international flights. Seven of the operators taking domestic reservations and 3 of the operators taking international reservations are female. If an operator is chosen at random, what is the probability that the person chosen takes domestic reservations or is a male? **inclusive; $\frac{10}{13}$**



- 18. MUSIC** Forty senior citizens were surveyed about their music preferences. The results are displayed in the Venn diagram. If a senior citizen from the survey group is selected at random, what is the probability that he or she likes only country and western music? What is the probability that he or she likes classical and/or country, but not 1940's pop?
- $\frac{3}{20}$; $\frac{2}{5}$**

- 2. TRAVEL** John is randomly selected to be given a chance to win a new car. He must choose a red or yellow marble from a bag containing 1 red, 2 yellow, 10 green, and 12 blue marbles. What is the probability he will win the car?
- $\frac{3}{25}$**

- 3. DICE** Alexis rolls two identical dice. What is the probability that the sum of the numbers rolled is odd? What is the probability that the sum of the numbers rolled is greater than 7? What is the probability that the sum of the numbers rolled is odd or greater than 7?
- $P(\text{odd}) = \frac{1}{2}$; $P(>7) = \frac{5}{12}$;**
 $P(\text{odd or } >7) = \frac{3}{4}$

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Chapter 12

Chapter 12

Glencoe Algebra 2

Chapter 12

39

Glencoe Algebra 2

NAME _____

DATE _____

PERIOD _____

12-6 Lesson Reading Guide

Statistical Measures

Get Ready for the Lesson

Read the introduction to Lesson 12-6 in your textbook.

There is more than one way to give an “average” score for this test. Three measures of central tendency for these scores are 94, 76.5 and 73.9. Can you tell which of these is the mean, the median, and the mode without doing any calculations? Explain your answer.

Sample answer: Yes. The mode must be one of the scores, so it must be an integer. The median must be either one of the scores or halfway between two of the scores, so it must be an integer or a decimal ending with .5. Therefore, 94 is the mode, 76.5 is the median, and 73.9 is the mean.

Read the Lesson

1. Match each measure with one of the six descriptions of how to find measures of central tendency and variation.

- a. median **vi**
- b. mode **i**
- c. range **iv**
- d. variance **iii**
- e. mean **ii**
- f. standard deviation **v**

- i. Find the most commonly occurring values or values in a set of data.
- ii. Add the data and divide by the number of items.
- iii. Find the mean of the squares of the differences between each value in the set of data and the mean.
- iv. Find the difference between the largest and smallest values in the set of data.
- v. Take the positive square root of the variance.
- vi. If there is an odd number of items in a set of data, take the middle one. If there is an even number of items, add the two middle items and divide by 2.

Remember What You Learned

2. It is usually easier to remember a complicated procedure if you break it down into steps. Write the procedure for finding the standard deviation for a set of data in a series of brief, numbered steps.

- Sample answer:**
1. Find the mean.
 2. Find the difference between each value and the mean.
 3. Square each difference.
 4. Find the mean of the squares.
 5. Take the positive square root.

Chapter 12

42

Glencoe Algebra 2

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____

DATE _____

PERIOD _____

12-6 Study Guide and Intervention

Statistical Measures

Measures of Central Tendency

Measures of Central Tendency	Use	When
mean	the data are spread out and you want an average of values	
median	the data contain outliers	
mode	the data are tightly clustered around one or two values	

Example Find the mean, median, and mode of the following set of data: {42, 39, 35, 40, 38, 35, 45}.

To find the mean, add the values and divide by the number of values.

$$\text{mean} = \frac{42 + 39 + 35 + 40 + 38 + 35 + 45}{7} \approx 39.14.$$

To find the median, arrange the values in ascending or descending order and choose the middle value. (If there is an even number of values, find the mean of the two middle values.) In this case, the median is 39.

To find the mode, take the most common value. In this case, the mode is 35.

Exercises

Find the mean, median, and mode of each set of data. Round to the nearest hundredth, if necessary.

1. {238, 261, 245, 249, 255, 262, 241, 245} **249.5; 247; 245**
2. {9, 13, 8, 10, 11, 9, 12, 16, 10, 9} **10.7; 10; 9**
3. {120, 108, 145, 129, 102, 132, 134, 118, 108, 142} **123.8; 124.5; 108**
4. {68, 54, 73, 58, 63, 72, 65, 70, 61} **64.89; 65; no mode**
5. {34, 49, 42, 38, 40, 45, 34, 28, 43, 30} **38.3; 39; 34**

6. The table at the right shows the populations of the six New England capitals. Which would be the most appropriate measure of central tendency to represent the data? Explain why and find that value. **There is no mode. The population of Boston is an outlier and would raise the mean too high. The median, 79,500, would be the best choice.**

City	Population (rounded to the nearest 1000)
Augusta, ME	19,000
Boston, MA	589,000
Concord, NH	37,000
Hartford, CT	122,000
Montpelier, VT	8,000
Providence, RI	174,000

Source: www.factfinder.census.gov

Chapter 12

43

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-6 Study Guide and Intervention *(continued)*

Statistical Measures

Measures of Variation The *range* and the **standard deviation** measure how scattered a set of data is.

Standard Deviation If a set of data consists of the n values x_1, x_2, \dots, x_n and has mean \bar{x} , then the standard deviation is given by $\sigma = \sqrt{\frac{(x_1 - \bar{x})^2 + (x_2 - \bar{x})^2 + \dots + (x_n - \bar{x})^2}{n}}$.

The square of the standard deviation is called the **variance**.

Example Find the variance and standard deviation of the data set {10, 9, 6, 9, 18, 4, 8, 20}.

Step 1 Find the mean.

$$\bar{x} = \frac{10 + 9 + 6 + 9 + 18 + 4 + 8 + 20}{8} = 10.5$$

Step 2 Find the variance.

$$\begin{aligned} \sigma^2 &= \frac{(x_1 - \bar{x})^2 + (x_2 - \bar{x})^2 + \dots + (x_n - \bar{x})^2}{n} && \text{Standard variance formula} \\ &= \frac{(10 - 10.5)^2 + (9 - 10.5)^2 + \dots + (20 - 10.5)^2}{8} \\ &= \frac{220}{8} \text{ or } 27.5 \end{aligned}$$

Step 3 Find the standard deviation.

$$\begin{aligned} \sigma &= \sqrt{27.5} \\ &\approx 5.2 \end{aligned}$$

The variance is 27.5 and the standard deviation is about 5.2.

Exercises

Find the variance and standard deviation of each set of data. Round to the nearest tenth.

- {100, 89, 112, 104, 96, 108, 93} **58.5; 7.6**
- {62, 54, 49, 62, 48, 53, 50} **29.4; 5.4**
- {8, 9, 8, 8, 9, 7, 8, 9, 6} **0.9; 0.9**
- {4.2, 5.0, 4.7, 4.5, 5.2, 4.8, 4.6, 5.1} **0.1; 0.3**

5. The table at the right lists the prices of ten brands of breakfast cereal. What is the standard deviation of the values to the nearest penny? **\$0.33**

Price of 10 Brands of Breakfast Cereal
\$2.29
\$3.19
\$3.39
\$2.79
\$2.99
\$3.09
\$3.19
\$2.59
\$2.79
\$3.29

NAME _____ DATE _____ PERIOD _____

12-6 Skills Practice

Statistical Measures

Find the variance and standard deviation of each set of data to the nearest tenth.

- {32, 41, 35, 35, 46, 42} **236.9, 4.9**
- {13, 62, 77, 24, 38, 19, 88} **763.8, 27.6**
- {89, 99, 42, 16, 42, 71, 16} **959.1, 31.0**
- {450, 400, 625, 225, 300, 750, 650, 625} **30,537.1; 174.7**
- {17, 23, 65, 94, 33, 33, 33, 8, 57, 75, 44, 12, 11, 68, 39} **630.7, 25.1**
- {7.2, 3.1, 3.8, 9.5, 8.3, 8.4} **5.8, 2.4**
- {1.5, 2.5, 3.5, 4.5, 4.5, 5.5, 6.5, 7.5} **3.5, 1.9**

For Exercises 8 and 9, use the table that shows the profit in billions of dollars reported by U.S. manufacturers for the first quarter of the years from 1997 through 2001.

Year	1997	1998	1999	2000	2001
Seasonally-Adjusted Profit (\$ billions)	\$61.4	\$75.6	\$60.9	\$78.5	\$45.3

Source: U.S. Census Bureau

- Find the mean and median of the data to the nearest tenth. **\$64.3 billion, \$61.4 billion**
- Which measure of central tendency best represents the data? Explain. **The median is more representative because the value 45.3 is not close to the other data points, and it lowers the mean.**

For Exercises 10 and 11, use the table that shows the percent of fourth grade students reading at or above the proficiency level in a nationally-administered reading assessment.

Year	1992	1994	1998	2000
Percent at or above proficiency level	29%	30%	31%	32%

Source: National Center for Education Statistics

- Find the mean, median, and standard deviation of the data to the nearest tenth. **30.5%, 30.5%, 1.1%**
- What do the statistics from Exercise 11 tell you about the data? **Sample answer: Since the median and mean are equal and the standard deviation is small, the percent of students reading at or above the proficiency level has not varied much from 1992 to 2000.**

NAME _____

DATE _____

PERIOD _____

12-6 Practice

Statistical Measures

Find the variance and standard deviation of each set of data to the nearest tenth.

- (47, 61, 93, 22, 82, 22, 37)
673.1, 25.9
- (10, 10, 54, 39, 96, 91, 91, 18)
1228.6, 35.1
- (1, 2, 2, 3, 3, 3, 4, 4, 4, 4, 5, 5, 5, 5)
1.6, 1.2
- (1100, 725, 850, 335, 700, 800, 950)
49,150.0; 221.7
- (3.4, 7.1, 8.5, 5.1, 4.7, 6.3, 9.9, 8.4, 3.6)
2.8, 0.5, 1.9, 0.8, 1.9, 1.5, 3.3, 2.6, 0.7, 2.5
0.8, 0.9

7. HEALTH CARE Eight physicians with 15 patients on a hospital floor see these patients an average of 18 minutes a day. The 22 nurses on the same floor see the patients an average of 3 hours a day. As a hospital administrator, would you quote the mean, median, or mode as an indicator of the amount of daily medical attention the patients on this floor receive? Explain. **Either the median or the mode; they are equal and higher than the mean, which is lowered by the smaller amount of time the physicians spend with the patients.**

For Exercises 8–10, use the frequency table that shows the percent of public school teachers in the U. S. in 1999 who used computers or the Internet at school for various administrative and teaching activities.

Activity	Percent Using Computer or Internet
Create instructional materials	39
Administrative record keeping	34
Communicate with colleagues	23
Gather information for planning lessons	16
Multimedia classroom presentations	8
Access research and best practices for teaching	8
Communicate with parents or students	8
Access model lesson plans	6

Source: National Assessment of Educational Progress

- Find the mean, median, and mode of the data. **17.75%, 12%, 8%**
- Suppose you believe teachers use computers or the Internet too infrequently. Which measure would you quote as the “average?” Explain. **Mode; it is lowest.**
- Suppose you believe teachers use computers or the Internet too often. Which measure would you quote as the “average?” Explain. **Mean; it is highest.**

For Exercises 11 and 12, use the frequency table that shows the number of games played by 24 American League baseball players between opening day, 2001 and September 8, 2001.

No. of Games	Frequency
141	4
140	3
139	4
138	5
137	2
136	3
135	3

Source: Major League Baseball

- Find the mean, median, mode, and standard deviation of the number of games played to the nearest tenth.
138.2, 138; 138, 2.0
- For how many players is the number of games within one standard deviation of the mean? **14**

Chapter 12

46

Glencoe Algebra 2

NAME _____

DATE _____

PERIOD _____

12-6 Word Problem Practice

Statistical Measures

1. SPORTS The table below shows the number of times some teams in the National Football League have won the Super Bowl.

NFL Team	Number of Super Bowl Victories
New England	3
Baltimore	2
Kansas City	1
St. Louis	1
Denver	2
Green Bay	1
Dallas	5
San Francisco	5
Oakland	2
Pittsburgh	5
Miami	2
Washington	3
NY Giants	2
NY Jets	1
Chicago	1

Source: www.pdqhelp.com

Which statistical measure represents the team(s) with the least Super Bowl victories? **the mode**

2. SALARIES The median salary in a small company is \$10.20 per hour. What percentage of the employees at the company earns more than \$10.20 per hour? **50%**

3. RANDOM GENERATORS Samuel has written a computer program to generate a random selection of the following two-digit numbers.

25, 67, 54, 99, 41, 87, 90, 18, 32

Find the mean, median, and mode of this data. **57; 54; none**

4. HEIGHTS The following table lists the heights of some of the great NBA players.

Player	Height (in inches)
Kareem Abdul-Jabbar	86
Larry Bird	81
Shaquille O’Neal	85
Wilt Chamberlain	85
Michael Jordan	78

Source: www.sivell.edu

Find the mean and standard deviation of the data in the table. Round your answer to the nearest hundredth.
83; 3.0

METEORS For Exercises 5–8, use the following information.

Arlene stayed up late one night to watch the Perseid meteor shower. She recorded the number of meteors she saw every ten minutes starting at 1 A.M. and going until 4 A.M. Her data are shown below.

8, 7, 8, 12, 17, 15, 22, 28, 29, 31, 28, 23, 29, 28, 25, 23, 15, 12

- What is the mean of this data set? **20**
- What is the median of this data set? **22.5**
- What is the mode of this data set? **28**
- What is the standard deviation of this data set? Round your answer to the nearest hundredth.
8.05

47

Chapter 12

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-6 Enrichment

Standard Deviation of Sample Data

A *population* is the set of all measurements of interest to an investigator. A *sample* is a subset of measurements selected from the population of interest. A *statistic* is any quantity whose value can be calculated from sample data. A common mistake is to use the terms *probability* and *statistics* interchangeably. Probabilities are used to make statements from a population to a sample, but statistics are calculated from a sample and are to make inferences about a population.

The *range* is a statistic calculated by taking the difference between the largest observation and the smallest observation. $\text{Range} = x_{\text{max}} - x_{\text{min}}$.

The *sample variance* is calculated using the formula, $s^2 = \frac{\sum_{i=1}^n (\bar{x} - x_i)^2}{n - 1}$ where \bar{x} is the sample mean. Therefore, the *sample standard deviation* is the square root of the sample variance, $s = \sqrt{s^2}$.

To calculate the sample variance:

1. Calculate the sample mean. For example, suppose a sample contains the numbers {2, 5, 6, 9, 11}. The sample mean is $\bar{x} = \frac{2 + 5 + 6 + 9 + 11}{5} = 6.6$.
2. Next use the formula above to calculate the sample variance, in this case:

$$s^2 = \frac{(6.6 - 2)^2 + (6.6 - 5)^2 + (6.6 - 6)^2 + (6.6 - 9)^2 + (6.6 - 11)^2}{4} = 12.3.$$
3. Finally, the sample standard deviation is equal to 3.507 by taking the square root of 12.3.

Exercises

1. What are some differences in the formula for the sample variance compared to the formula for the population variance? **It uses the sample mean instead of the population mean, and since the sample mean is an estimator for the population mean, the denominator is $n - 1$ instead of n .**
2. Given the random sample {5, 7, 1, 2, 4}, find the sample variance. **5.70**
3. Calculate the sample standard deviation. **2.587**
4. Calculate the range of the sample data {5, 7, 1, 2, 4}. **6**
5. An approximation for the sample standard deviation is given by: $s \approx \frac{\text{Range}}{4}$. Compare this answer to your answer from 3. **6**

NAME _____ DATE _____ PERIOD _____

12-7 Reading to Learn Mathematics

The Normal Distribution

Get Ready for the Lesson

Read the introduction to Lesson 12-7 in your textbook.

There were 66 players on the team and the mean height was approximately 74.1. About what fraction of the players' heights are between 72 and 75, inclusive?

Sample answer: about $\frac{1}{2}$

Read the Lesson

1. Indicate whether each of the following statements is *true* or *false*.
 - a. In a continuous probability distribution, there is a finite number of possible outcomes. **false**
 - b. Every normal distribution can be represented by a bell curve. **true**
 - c. A distribution that is represented by a curve that is high at the left and has a tail to the right is negatively skewed. **false**
 - d. A normal distribution is an example of a skewed distribution. **false**

2. Ms. Rose gave the same quiz to her two geometry classes. She recorded the following scores.

First-period class:

Score	0	1	2	3	4	5	6	7	8	9	10
Frequency	1	0	1	0	3	4	5	7	4	3	2

Fifth-period class:

Score	0	1	2	3	4	5	6	7	8	9	10
Frequency	0	0	0	0	3	4	9	7	6	1	0

In each class, 30 students took the quiz. The mean score for each class was 6.4. Which set of scores has the greater standard deviation? (Answer this question without doing any calculations.) Explain your answer.

First period class; sample answer: The scores are more spread out from the mean than for the fifth period class.

Remember What You Learned


3. Many students have trouble remembering how to determine if a curve represents a distribution that is *positively skewed* or *negatively skewed*. What is an easy way to remember this?

Sample answer: Follow the tail! If the tail is on the right (positive direction), the distribution is positively skewed. If the tail is on the left (negative direction), the distribution is negatively skewed.

12-7 Study Guide and Intervention (continued)

The Normal Distribution

Use Normal Distributions




Normal distributions have these properties.
The graph is maximized at the mean.
The mean, median, and mode are about equal.
About 68% of the values are within one standard deviation of the mean.
About 95% of the values are within two standard deviations of the mean.
About 99% of the values are within three standard deviations of the mean.

LESSON 12-7

Example The heights of players in a basketball league are normally distributed with a mean of 6 feet 1 inch and a standard deviation of 2 inches.

a. What is the probability that a player selected at random will be shorter than 5 feet 9 inches?

Draw a normal curve. Label the mean and the mean plus or minus multiples of the standard deviation.
The value of 5 feet 9 inches is 2 standard deviations below the mean, so approximately 2.5% of the players will be shorter than 5 feet 9 inches.



b. If there are 240 players in the league, about how many players are taller than 6 feet 3 inches?

The value of 6 feet 3 inches is one standard deviation above the mean. Approximately 16% of the players will be taller than this height.
 $240 \times 0.16 \approx 38$
About 38 of the players are taller than 6 feet 3 inches.

Exercises

EGG PRODUCTION The number of eggs laid per year by a particular breed of chicken is normally distributed with a mean of 225 and a standard deviation of 10 eggs.

- About what percent of the chickens will lay between 215 and 235 eggs per year? **68%**
- In a flock of 400 chickens, about how many would you expect to lay more than 245 eggs per year? **10 chickens**

MANUFACTURING The diameter of bolts produced by a manufacturing plant is normally distributed with a mean of 18 mm and a standard deviation of 0.2 mm.


- What percent of bolts coming off of the assembly line have a diameter greater than 18.4 mm? **2.5%**
- What percent have a diameter between 17.8 and 18.2 mm? **68%**

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

12-7 Study Guide and Intervention

The Normal Distribution

Normal and Skewed Distributions A continuous probability distribution is represented by a curve.



Example Determine whether the data below appear to be *positively skewed*, *negatively skewed*, or *normally distributed*.
{100, 120, 110, 100, 110, 80, 100, 90, 100, 120, 100, 90, 110, 100, 90, 80, 100, 90}

Make a frequency table for the data.

Value	80	90	100	110	120
Frequency	2	4	7	3	2


Then use the data to make a histogram.
Since the histogram is roughly symmetric, the data appear to be normally distributed.

Exercises

Determine whether the data in each table appear to be positively skewed, negatively skewed, or normally distributed. Make a histogram of the data.


- | | | | | | | | | | | | | | | |
|-----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| 127 | 24 | 29 | 25 | 27 | 22 | 24 | 25 | 22 | 27 | 24 | 22 | 25 | 24 | 22 |
|-----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|

positively skewed




- | Shoe Size | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|-----------------|---|---|---|---|---|---|----|
| No. of Students | 1 | 2 | 4 | 8 | 5 | 1 | 2 |

normally distributed



- | Housing Price | No. of Houses Sold |
|---------------------|--------------------|
| less than \$100,000 | 0 |
| \$100,000–\$120,000 | 1 |
| \$121,000–\$140,000 | 3 |
| \$141,000–\$160,000 | 7 |
| \$161,000–\$180,000 | 8 |
| \$181,000–\$200,000 | 6 |
| over \$200,000 | 12 |

negatively skewed



NAME _____ DATE _____ PERIOD _____

12-7 Skills Practice

The Normal Distribution

Determine whether the data in each table appear to be *positively skewed*, *negatively skewed*, or *normally distributed*.

1.

Miles Run	Track Team Members
0–4	3
5–9	4
10–14	7
15–19	5
20–23	2

2.


Speeches Given	Political Candidates
0–5	1
6–11	2
12–17	3
18–23	8
24–29	8

normally distributed

negatively skewed

For Exercises 3 and 4, use the frequency table that shows the average number of days patients spent on the surgical ward of a hospital last year.

Days	Number of Patients
0–3	5
4–7	18
8–11	11
12–15	9
16+	6



3. Make a histogram of the data.
4. Do the data appear to be *positively skewed*, *negatively skewed*, or *normally distributed*? Explain.
Positively skewed; the histogram is high at the left and has a tail to the right.

DELIVERY For Exercises 5–7, use the following information.

- The time it takes a bicycle courier to deliver a parcel to his farthest customer is normally distributed with a mean of 40 minutes and a standard deviation of 4 minutes.
5. About what percent of the courier's trips to this customer take between 36 and 44 minutes?
68%
6. About what percent of the courier's trips to this customer take between 40 and 48 minutes?
47.5%
7. About what percent of the courier's trips to this customer take less than 32 minutes?
2.5%

TESTING For Exercises 8–10, use the following information.

- The average time it takes sophomores to complete a math test is normally distributed with a mean of 63.3 minutes and a standard deviation of 12.3 minutes.
8. About what percent of the sophomores take more than 75.6 minutes to complete the test?
16%
9. About what percent of the sophomores take between 51 and 63.3 minutes?
34%
10. About what percent of the sophomores take less than 63.3 minutes to complete the test?
50%

NAME _____ DATE _____ PERIOD _____

12-7 Practice

The Normal Distribution

Determine whether the data in each table appear to be *positively skewed*, *negatively skewed*, or *normally distributed*.

1.

Minutes	Frequency
0–25	27
26–50	46
51–75	89
75–100	57
100+	24

2.


Average Age of High School Principals	Number
31–35	3
36–40	8
41–45	15
46–50	32
51–55	40
56–60	38
60+	4

normally distributed

negatively skewed

For Exercises 3 and 4, use the frequency table that shows the number of hours worked per week by 100 high school seniors.

Hours	Number of Students
0–8	30
9–17	45
18–25	20
26+	5



3. Make a histogram of the data.
4. Do the data appear to be *positively skewed*, *negatively skewed*, or *normally distributed*? Explain.
Positively skewed; the histogram is high at the left and has a tail to the right.

TESTING For Exercises 5–10, use the following information.

- The scores on a test administered to prospective employees are normally distributed with a mean of 100 and a standard deviation of 15.
5. About what percent of the scores are between 70 and 130?
95%
6. About what percent of the scores are between 85 and 130?
81.5%
7. About what percent of the scores are over 115?
16%
8. About what percent of the scores are lower than 85 or higher than 115?
32%
9. If 80 people take the test, how many would you expect to score higher than 130?
2
10. If 75 people take the test, how many would you expect to score lower than 85?
12


11. TEMPERATURE The daily July surface temperature of a lake at a resort has a mean of 82° and a standard deviation of 4.2°. If you prefer to swim when the temperature is at least 77.8°, about what percent of the days does the temperature meet your preference?
84%

12-7 Word Problem Practice

The Normal Distribution

1. PARKING Over several years, Bertram conducted a study of how far into parking spaces people tend to park by measuring the distance from the end of a parking space to the front fender of a car parked in the space. He discovered that the distribution of the data closely approximated a normal distribution with mean 8.5 inches. He found that about 5% of cars parked more than 11.5 inches away from the end of the parking space. What percentage of cars would you expect parked less than 5.5 inches away from the end of the parking space?
5%

2. HEIGHT Chandra's graph of the number of tenth grade students of different heights is shown below.



Is the data positively skewed, negatively skewed, or normally distributed?
Negatively skewed

3. OVENS An oven manufacturer tries to make the temperature setting on its ovens as accurate as possible. However, if one measures the actual temperatures in the ovens when the temperature setting is 350°F, they will differ slightly from 350°F. The set of actual temperatures for all the ovens is normally distributed around 350°F with a standard deviation of 0.5°F. About what percentage of ovens will be between 350°F and 351°F when their temperature setting is 350°F?
47.5%

12-7 Enrichment

Calculating Z-Scores

The normal distribution is the most important probability distribution. Many physical measurements have distributions approximately normal. Examples include height, weight, and measures of intelligence. More importantly, even if the individual variables are not normally distributed, sums and averages tend to still be normally distributed. Unfortunately, normal probability distribution functions are difficult to calculate. Fortunately, statisticians have compiled a table for a normal distribution with mean of zero and standard deviation of one. This is called the Standard Normal Distribution and is typically denoted by $N(0, 1)$, where the N indicates a normal distribution which has mean, μ ($\mu = 0$), and standard deviation, σ ($\sigma = 1$).

Suppose the variable x is normally distributed with mean μ and standard deviation σ . In order to calculate probabilities of this normal distribution, we must standardize the variable x by an appropriate transformation. The letter Z denotes the transformed variable and is called the Z -score, which is a measure of relative standing. The following steps are needed to complete the transformation.

- If the mean and standard deviation are not given, then calculate the mean and standard deviation of the given (population) data.
- Define $Z = \frac{x - \mu}{\sigma}$.

Example Find the standard normal variable Z given and $\mu = 15$ and $\sigma = 3$.

Apply the transform to the variable X using the definition above, that is: $Z = \frac{X - 15}{3}$.

- 1.** Suppose that the time, X , to complete an exam is normally distributed. The time, in minutes, of a class of 12 to complete the exam is given in the table. Transform X to a Z -score.

Student	1	2	3	4	5	6	7	8	9	10	11	12
Time	35	42	48	33	32	39	40	52	48	34	36	44

Mean = 40.25, Standard Deviation = 6.34
 $Z = (X - 40.25) / 6.34$

- 2.** Suppose that a random variable X is normally distributed with $\mu = 20$ and $\sigma = 5$. Convert the following probability statements to the equivalent statements by standardizing X .

Example $P(X < 25) = P\left(\frac{X - 20}{5} < 25\right) = P(Z < 25)$

- a. $P(X > 18)$
 $P\left(Z > \frac{18 - 20}{5}\right) = P\left(Z > -\frac{2}{5}\right)$
- b. $P(17 < X < 23)$
 $P\left(\frac{17 - 20}{5} < Z < \frac{23 - 20}{5}\right) = P\left(-\frac{3}{5} < Z < \frac{3}{5}\right)$
- c. $P(X < 19)$
 $P\left(Z < \frac{19 - 20}{5}\right) = P\left(Z < -\frac{1}{5}\right)$

NAME _____ DATE _____ PERIOD _____

12-8 Lesson Reading Guide

Exponential and Binomial Distribution

Get Ready for the Lesson

Read the introduction to Lesson 12-8 in your textbook.

Is a randomly chosen student likely to be one that talks on the phone for a very short period of time or for a very long period of time? **No, most people will talk an average amount of time.**

Read the Lesson

- Indicate whether each situation can be represented using an exponential distribution or a binomial distribution.
 - You are trying to predict how many times a coin will land with the tails side up if you flip it 50 times. **binomial distribution**
 - You would like to find the probability that there will be more than 5 pink gumballs in a bag of assorted color gumballs. **binomial distribution**
 - You are trying to predict how long your refrigerator will last. **exponential distribution**
 - You are calculating the probability that a person in your class will be taller than 5 feet, 5 inches. **binomial distribution**
 - You would like to determine the percentage of cellular phones that will last longer than 7 years and the percentage that will last longer than 5 years. **exponential distribution**
 - You want to predict the probability that a person in your neighborhood is older than you are. **binomial distribution**
- Write an equation that can be used to calculate each of the following:
 - The expected number of successes in a binomial distribution that has a 30% rate of success when there are 50 trials. **$E = (50)(0.3)$**
 - The probability that a randomly selected number from an exponential distribution will be greater than 4 if the mean is 1.5. **$P = e^{-\frac{2}{3}(4)}$**

Remember What You Learned

- In binomial distributions, the only possible outcomes are success and failure, but sometimes binomial experiments include events that can have several results. Explain how this is possible. **For example, in 1b, there are assorted colors of gumballs, but you are concerned about how many pink ones there will be. For evaluation as a binomial distribution, a "success" is pink. Any other color is considered a "failure".**

Chapter 12

56

Glencoe Algebra 2

NAME _____ DATE _____ PERIOD _____

12-8 Study Guide and Intervention

Exponential and Binomial Distribution

Exponential Distribution


Exponential Distributions are used to predict the probabilities of events based on time.

Probability that a randomly chosen domain value for the exponential function $f(x)$ will be greater than the given value of x . The value m is the multiplicative inverse of the mean.	$f(x) = e^{-mx}$
Probability that a randomly chosen domain value for the exponential function $f(x)$ will be less than the given value of x . The value m is the multiplicative inverse of the mean.	$f(x) = 1 - e^{-mx}$

Example An exponential distribution function has a mean of 2. Graph the distribution function and label the mean. What is the probability that a randomly chosen value of x will be less than 3?

The equation for the function will be $f(x) = e^{-0.5x}$, where m is the multiplicative inverse of the mean. Since the mean is 2, the value of m will be $\frac{1}{2}$ or 0.5. Substituting the value of m into the equation for an exponential distribution, $f(x) = e^{-0.5x}$.

Since the function is applicable when x is greater than zero, the graph only includes the first quadrant x and y values. The y -axis represents the probability, which ranges from 0 to 1. The probability that a randomly chosen value will be less than 3 can be found using the graph or the equation for the distribution. From the graph, the value of $f(x)$ is about 0.22 when x is 3, which means that the probability that x is greater than 3 is 0.22. Remember to subtract the value from one when you want to know the probability that a randomly chosen value will be less than the given value. The probability that x will be less than 3 is $1 - 0.22$, or 0.88.



To use the equation developed for the function to find the probability, substitute 3 for the value of x into the equation $f(x) = 1 - e^{-0.5x}$ and solve. The probability that x will be greater than 3 is 0.88.

Exercises

- Write the equation for an exponential distribution that has a mean of 0.5. **$f(x) = e^{-2x}$**
- An exponential distribution has a mean of 6. Find each probability.
- $x > 7$ **0.31**
 - $x > 10$ **0.19**
 - $x > 2$ **0.72**
 - $x < 9$ **0.78**
 - $x < 7$ **0.69**
 - $x < 3$ **0.88**

Chapter 12

57

Glencoe Algebra 2

12-8 Study Guide and Intervention**Exponential and Binomial Distribution****Binomial Distribution**

Binomial distributions occur when a series of trials only has two possible outcomes, success and failure, and the probability for success is the same in each trial.

Probability of x successes in n independent trials	$P(x) = C(n, x) p^x q^{n-x}$ where p is the probability of success of an individual trial and q is the probability of failure on that same individual trial ($p + q = 1$).
Expectation of a Binomial Distribution	$E(x) = np$ where n is the total number of trials and p is the probability of success.

Example You are going to flip a coin 10 times. What is the probability that 3 of the coin flips will result in heads?

A success in this case would be heads, and there is an equal chance that a flip will result in heads or tails, so the probability of success is 0.5.

$$P(x) = C(n, x) p^x q^{n-x}$$

Probability Formula

$$P(x) = C(10, 3) (0.5)^3 (0.5)^{10-3}$$

$n = 10, x = 3, p = 0.5$

$$P(x) = \frac{10!}{(10-3)! 3!} (0.5)^3 (0.5)^{10-3}$$

Combination Formula

$$P(x) = 120(0.5)^3(0.5)^7 \text{ or } 0.12$$

Simplify.

The probability of getting 3 heads is 0.12.

Exercises

A binomial distribution has a 30% rate of success. There are 20 trials. Find each probability.

1. 10 successes **0.03** 2. 3 successes **0.07** 3. 6 successes **0.19** 4. 15 failures **0.18**

The probability of getting a red fruit candy from an assorted bag is 0.1. There are 50 pieces of candy in a bag.

5. What is the probability that there are at least 4 pieces of red candy in the bag? **0.75**
6. What is the probability that there are 8 pieces of red candy in the bag? **0.06**
7. What is the probability that there are 2 or less pieces of red candy in the bag? **0.11**
8. What is the expected number of red pieces of candy in the bag? **5**


12-8 Skills Practice**Exponential and Binomial Distribution**

For Exercises 1–4, use the following information. The exponential distribution shown at the right has a mean of 3. Use the graph or function equation to find each probability. Round to the nearest tenth.

1. $x > 7$ **0.1**
2. $x < 8$ **0.9**
3. $x < 5$ **0.8**
4. $x > 2$ **0.5**

For Exercises 5–6, use the following information. A binomial distribution has a 80% rate of success. There are 65 trials.

5. What is the expected number of successes? **52**
6. What is the expected number of failures? **13**



For Exercises 7–11, use the following information. A binomial distribution has a 40% rate of success. There are 23 trials.

7. What is the probability that there will be 13 successes? **0.05**
8. What is the probability that there will be 15 failures? **0.15**
9. What is the probability that there will be at least 4 successes? **0.99**
10. What is the probability that there will be 20 or more failures? **0.005**
11. What is the expected number of successes? **about 9**

MAIL For Exercises 12–14, use the following information.

The amount of time it takes a parcel of mail to travel first class from San Francisco to New York follows an exponential distribution. The mean travel time is 5 days.

12. What is the probability that a randomly selected package will take longer than 7 days? **0.25**
13. What is the probability that a randomly selected package will take less than 6 days? **0.70**
14. What is the probability that a randomly selected package will take longer than 3 days? **0.55**
- SOCKS For Exercises 15–17, use the following information.**
You grab 6 pairs of socks out of your drawer for a trip without looking. The socks are 75% white.
15. What is the probability that 5 pairs of the socks are white? **0.36**
16. What is the probability that at least 3 pairs of the socks are white? **0.97**
17. What is the probability that all of the socks are white? **0.18**

NAME _____

DATE _____

PERIOD _____

12-8 Practice

Exponential and Binomial Distribution

For Exercises 1–4, use the following information. The exponential distribution shown at the right has a mean of 0.25. Use the graph or function equation to find each probability. Round to the nearest hundredth.

- $x > 0.6$ **0.09**
- $x > 1$ **0.02**
- $x < 0.75$ **0.95**
- $x < 0.5$ **0.86**

For Exercises 5–9, use the following information. A binomial distribution has a 25% rate of success. There are 41 trials.

- What is the probability that there will be 11 successes? **0.13**
- What is the probability that there will be 26 failures? **0.03**
- What is the probability that there will be at least 3 successes? **0.999**
- What is the probability that there will be 38 or more failures? **0.004**
- What is the expected number of failures? **about 31**

HYGIENE For Exercises 10–13, use the following information.

- The mean amount of time a person spends brushing his or her teeth is 45 seconds.
- What is the probability that a randomly selected person spends more than 2 minutes brushing his or her teeth? **0.07**
 - What is the probability that a randomly selected person spends less than 1.5 minutes brushing his or her teeth? **0.86**
 - 75% of people spend less than how long brushing their teeth? **62 seconds**
 - 10% of people spend more than how long brushing their teeth? **104 seconds**

PROJECTS For Exercises 14–17, use the following information.

- Your science teacher is drawing names from a hat to assign groups of 4 for a project. Half of the people in your class are girls.
- What is the expected number of girls in a group? **2**
 - What is the probability that there will be 3 boys in a group? **0.25**
 - What is the probability that there will be 2 or less girls in a group? **0.69**
 - What is the probability that all four members of a group will be boys? **0.0625**

Chapter 12

60

Glencoe Algebra 2

NAME _____

DATE _____

PERIOD _____

12-8 Word Problem Practice

Exponential and Binomial Distribution

1. CARS A specific brand of timing belt has a mean life of 75,000 miles. The manufacturer encourages customers to change their timing belts at 60,000; when the belts fail, they can damage the engine. Sam's car has 63,000 miles on it, but he is planning to buy a new car at the end of the summer. He thinks he will put about 3,000 more miles on the car over the summer. He is trying to decide whether to replace the timing belt now or save the money to spend when he buys a new car. What is the probability that Sam's timing belt will last longer than the mileage he predicted that he will have at the end of the summer? Based on the probability, should he replace the belt? **0.41, yes**

SPORTS For Exercises 4 and 5, use the following information.

Recent studies suggest that 13% of the population is left-handed. Your baseball team has 9 members

- What is the probability that at least one member of your team is left-handed? **0.71**

Lesson 12-8

- What is the probability that three members of your team are left-handed? **0.08**

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

2. HOME IMPROVEMENT The average life of the seals around a particular brand of window is 20 years. Penny is buying a home that is 12 years old and has the above-mentioned brand of windows. What is the probability that the seals are still effective? **0.55**

3. BUDGET The average 4-person family spends about \$140 per month on groceries. The probability is 0.51 that a family spends less than how much? Round to the nearest dollar. **\$100**

- DECORATING** Susan is on the decorating committee for the spring dance at her school. The committee has chosen to use strands of white lights as part of the decoration. The probability of having a defective light bulb is 0.002. Susan is using five strands of 500-lights each. What is the probability that at least 4 bulbs will be defective? **0.74**

Chapter 12

61

Glencoe Algebra 2


12-8 Enrichment

Exploring Binomial Distribution Functions

In Lesson 12-8, you calculated probabilities using binomial distributions. In this exercise, you will explore graphs of binomial distributions.


- Bob is the pitcher on your baseball team. The probability that he will throw a strike is 50%. If Bob throws six pitches, fill in the probabilities for each number of strikes in the table below. Then graph the number of strikes (s) versus probability (p) on the coordinate grid provided.

Strikes (s)	Probability (p)
0	0.016
1	0.09
2	0.23
3	0.31
4	0.23
5	0.09
6	0.016



- Joy's pug is going to have puppies. The probability that a puppy will be black is 25%. If there are 6 puppies in the litter, fill in the table below with the probability that corresponds to each possible number of black puppies. Then graph the number of black puppies (b) versus probability(p) on the coordinate grid provided.

Black Puppies (b)	Probability (p)
0	0.18
1	0.36
2	0.30
3	0.13
4	0.03
5	0.004
6	0.0002



- How do the graphs relate to the distribution curves that you studied in Lesson 12-7? **The graph in Exercise 1 is a normal distribution and the graph in Exercise 2 is a positively skewed distribution.**
- What general shape would you expect the graph of a binomial distribution with a 50% probability of success to look like? With a probability less than 50%? With a probability greater than 50%?



12-9 Lesson Reading Guide

Binomial Experiments

Get Ready for the Lesson

Read the introduction to Lesson 12-9 in your textbook.

Suppose you are taking a 50-question multiple-choice test in which there are 5 answer choices for each question. You are told that no points will be deducted for wrong answers. Should you guess the answers to the questions you do not know? Explain your reasoning.

Sample answer: Yes; the probability of guessing the right answer to a question is $\frac{1}{5}$, so you have a chance to get some points by guessing, and you have nothing to lose.

Read the Lesson

- Indicate whether each of the following is a *binomial experiment* or *not a binomial experiment*. If the experiment is not a binomial experiment, explain why.

- A fair coin is tossed 10 times and “heads” or “tails” is recorded each time. **binomial experiment**
- A pair of dice is thrown 5 times and the sum of the numbers that come up is recorded each time. **Not a binomial experiment; there are more than two possible outcomes for each trial.**
- There are 5 red marbles and 6 blue marbles in a bag. One marble is drawn from the bag and its color recorded. The marble is not put back in the bag. A second marble is drawn and its color recorded. **Not a binomial experiment; the trials are not independent (or, the probabilities for the two trials are not the same).**
- There are 5 red marbles and 6 blue marbles in a bag. One marble is drawn from the bag and its color recorded. The marble is put back in the bag. A second marble is drawn and its color recorded. **binomial experiment**

- Len randomly guesses the answers to all 6 multiple-choice questions on his chemistry test. Each question has 5 choices. Which of the following expressions gives the probability that he will get at least 4 of the answers correct? **B**

A. $P(6, 4) \left(\frac{1}{5}\right)^4 \left(\frac{4}{5}\right)^2 + P(6, 5) \left(\frac{1}{5}\right)^5 \left(\frac{4}{5}\right)^1 + P(6, 6) \left(\frac{1}{5}\right)^6 \left(\frac{4}{5}\right)^0$
 B. $C(6, 4) \left(\frac{1}{5}\right)^4 \left(\frac{4}{5}\right)^2 + C(6, 5) \left(\frac{1}{5}\right)^5 \left(\frac{4}{5}\right)^1 + C(6, 6) \left(\frac{1}{5}\right)^6 \left(\frac{4}{5}\right)^0$
 C. $C(6, 4) \left(\frac{1}{5}\right)^2 \left(\frac{4}{5}\right)^4 + C(6, 5) \left(\frac{1}{5}\right)^1 \left(\frac{4}{5}\right)^5 + C(6, 6) \left(\frac{1}{5}\right)^0 \left(\frac{4}{5}\right)^6$

Remember What You Learned

- Some students have trouble remembering how to calculate binomial probabilities. What is an easy way to remember which numbers to put into an expression like $C(6, 4) \left(\frac{1}{5}\right)^4 \left(\frac{4}{5}\right)^2$? **Sample answer: The binomial coefficient is $C(n, r)$, where n is the number of trials and r is the number of successes. The probability of success is raised to the r th power and the probability of failure is raised to the $(n - r)$ th power.**

NAME _____ DATE _____ PERIOD _____

12-9 Study Guide and Intervention

Binomial Experiments

Binomial Expansions For situations with only 2 possible outcomes, you can use the Binomial Theorem to find probabilities. The coefficients of terms in a binomial expansion can be found by using combinations.

Example What is the probability that 3 coins show heads and 3 show tails when 6 coins are tossed?

There are 2 possible outcomes that are equally likely: heads (H) and tails (T). The tosses of 6 coins are independent events. When $(H + T)^6$ is expanded, the term containing H^3T^3 , which represents 3 heads and 3 tails, is used to get the desired probability. By the Binomial Theorem the coefficient of H^3T^3 is $C(6, 3)$.

$$P(3 \text{ heads, } 3 \text{ tails}) = \frac{6!}{3!3!} \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^3 \quad P(H) = \frac{1}{2} \text{ and } P(T) = \frac{1}{2}$$

$$= \frac{20}{64}$$

$$= \frac{5}{16}$$

The probability of getting 3 heads and 3 tails is $\frac{5}{16}$ or 0.3125.

Exercises

Find each probability if a coin is tossed 8 times.

- $P(\text{exactly } 5 \text{ heads})$
about 22%
- $P(\text{exactly } 2 \text{ heads})$
about 11%
- $P(\text{even number of heads})$
50%
- $P(\text{at least } 6 \text{ heads})$
about 14%

Mike guesses on all 10 questions of a true-false test. If the answers true and false are evenly distributed, find each probability.

- Mike gets exactly 8 correct answers.
 $\frac{45}{1024}$ or 0.044
- Mike gets at most 3 correct answers.
 $\frac{11}{64}$ or 0.172

7. A die is tossed 4 times. What is the probability of tossing exactly two sixes?

$\frac{25}{216}$ or 0.116

NAME _____ DATE _____ PERIOD _____

12-9 Study Guide and Intervention

Binomial Experiments

A binomial experiment is possible if and only if all of these conditions occur.

- There are exactly two outcomes for each trial.
- There is a fixed number of trials.
- The trials are independent.
- The probabilities for each trial are the same.

Example Suppose a coin is weighted so that the probability of getting heads in any one toss is 90%. What is the probability of getting exactly 7 heads in 8 tosses?

The probability of getting heads is $\frac{9}{10}$, and the probability of getting tails is $\frac{1}{10}$. There are $C(8, 7)$ ways to choose the 7 heads.

$$P(7 \text{ heads}) = C(8, 7) \left(\frac{9}{10}\right)^7 \left(\frac{1}{10}\right)^1$$

$$= 8 \cdot \frac{9^7}{10^8}$$

$$\approx 0.38$$

The probability of getting 7 heads in 8 tosses is about 38%.

Exercises

1. BASKETBALL For any one foul shot, Derek has a probability of 0.72 of getting the shot in the basket. As part of a practice drill, he shoots 8 shots from the foul line.

- What is the probability that he gets in exactly 6 foul shots? **about 31%**
- What is the probability that he gets in at least 6 foul shots? **about 60%**

2. SCHOOL A teacher is trying to decide whether to have 4 or 5 choices per question on her multiple choice test. She wants to prevent students who just guess from scoring well on the test.

- On a 5-question multiple-choice test with 4 choices per question, what is the probability that a student can score at least 60% by guessing? **10.4%**
- What is the probability that a student can score at least 60% by guessing on a test of the same length with 5 choices per question? **5.8%**

3. Julie rolls two dice and adds the two numbers.

- What is the probability that the sum will be divisible by 3? **$\frac{1}{3}$**
- If she rolls the dice 5 times what is the chance that she will get exactly 3 sums that are divisible by 3? **about 16%**

4. SKATING During practice a skater falls 15% of the time when practicing a triple axel. During one practice session he attempts 20 triple axels.

- What is the probability that he will fall only once? **about 14%**
- What is the probability that he will fall 4 times? **about 18%**

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____ DATE _____ PERIOD _____

12-9 Practice

Binomial Experiments

Find each probability if a coin is tossed 6 times.

- $P(4 \text{ heads}) = \frac{1}{16}$
- $P(0 \text{ heads}) = \frac{1}{16}$
- $P(\text{exactly 3 heads}) = \frac{1}{4}$
- $P(\text{exactly 2 heads}) = \frac{3}{8}$
- $P(\text{exactly 1 head}) = \frac{1}{4}$
- $P(\text{at least 3 heads}) = \frac{5}{16}$

Find each probability if a die is rolled 3 times.

- $P(\text{exactly one 2}) = \frac{25}{72}$
- $P(\text{exactly two 2s}) = \frac{5}{72}$
- $P(\text{exactly three 2s}) = \frac{1}{216}$
- $P(\text{at most one 2}) = \frac{25}{27}$

A town that presents a fireworks display during its July 4 celebration found the probability that a family with two or more children will watch the fireworks is $\frac{3}{5}$. If 5 of these families are selected at random, find each probability.

- $P(\text{exactly 3 families watch the fireworks}) = \frac{216}{625}$
- $P(\text{exactly 2 families watch the fireworks}) = \frac{144}{625}$
- $P(\text{exactly 5 families watch the fireworks}) = \frac{243}{3125}$
- $P(\text{no families watch the fireworks}) = \frac{32}{3125}$
- $P(\text{at least 4 families watch the fireworks}) = \frac{1053}{3125}$
- $P(\text{at most 1 family watches the fireworks}) = \frac{272}{3125}$

One section of a standardized English language test has 10 true/false questions. Find each probability when a student guesses at all ten questions.

- $P(\text{exactly 8 correct}) = \frac{45}{1024}$
- $P(\text{exactly 2 correct}) = \frac{45}{1024}$
- $P(\text{exactly half correct}) = \frac{63}{256}$
- $P(\text{all 10 correct}) = \frac{1}{1024}$
- $P(0 \text{ correct}) = \frac{1}{1024}$
- $P(\text{at least 8 correct}) = \frac{7}{128}$

Chapter 12

Glencoe Algebra 2

66

NAME _____ DATE _____ PERIOD _____

12-9 Skills Practice

Binomial Experiments

Find each probability if a coin is tossed 4 times.

- $P(4 \text{ heads}) = \frac{1}{16}$
- $P(0 \text{ heads}) = \frac{1}{16}$
- $P(\text{exactly 3 heads}) = \frac{1}{4}$
- $P(\text{exactly 2 heads}) = \frac{3}{8}$
- $P(\text{exactly 1 head}) = \frac{1}{4}$
- $P(\text{at least 3 heads}) = \frac{5}{16}$

The probability of Chris making a free throw is $\frac{2}{3}$. If she shoots 5 times, find each probability.

- $P(\text{all missed}) = \frac{1}{243}$
- $P(\text{all made}) = \frac{32}{243}$
- $P(\text{exactly 2 made}) = \frac{40}{243}$
- $P(\text{exactly 1 missed}) = \frac{80}{243}$
- $P(\text{at least 3 made}) = \frac{64}{81}$
- $P(\text{at most 2 made}) = \frac{17}{81}$

When Tarin and Sam play a certain board game, the probability that Tarin will win a game is $\frac{3}{4}$. If they play 5 games, find each probability.

- $P(\text{Sam wins only once}) = \frac{405}{1024}$
- $P(\text{Sam wins exactly twice}) = \frac{45}{512}$
- $P(\text{Sam wins exactly 3 games}) = \frac{45}{512}$
- $P(\text{Sam wins at least 1 game}) = \frac{781}{1024}$
- $P(\text{Tarin wins at least 3 games}) = \frac{459}{512}$
- $P(\text{Tarin wins at most 2 games}) = \frac{53}{512}$

19. SAFETY In August 2001, the American Automobile Association reported that 73% of Americans use seat belts. In a random selection of 10 Americans in 2001, what is the probability that exactly half of them use seat belts? **about 7.5%**

HEALTH For Exercises 20 and 21, use the following information.

In 2001, the American Heart Association reported that 50 percent of the Americans who receive heart transplants are ages 50–64 and 20 percent are ages 35–49.

- In a randomly selected group of 10 heart transplant recipients, what is the probability that at least 8 of them are ages 50–64? $\frac{7}{128}$
- In a randomly selected group of 5 heart transplant recipients, what is the probability that 2 of them are ages 35–49? $\frac{128}{625}$

Chapter 12

Glencoe Algebra 2

67

NAME _____ DATE _____ PERIOD _____

12-9 Word Problem Practice

Binomial Experiments

- GENETICS** Dagmar is conducting a genetic experiment. Before she performs the experiment, she would like to compute theoretically probabilities for some of the outcomes. One of these computations involves expanding $(p + q)^4$. What is this expansion?
 $p^4 + 4p^3q + 6p^2q^2 + 4pq^3 + q^4$

- GAMES** The probability that Kendra will win a card game is $\frac{2}{3}$. If Kendra plays 7 games, what is the probability she wins exactly 4 games? Round your answer to the nearest thousandth.
about 0.256

- DEFECTS** An electronics parts manufacturer produces capacitors for electronic circuits. The probability that a capacitor comes out defective is 1 in 1,000. In a batch of 10,000 capacitors, write an expression for the probability that 10 of the capacitors are defective.
 $C(10000, 10) \left(\frac{1}{1000}\right)^{10} \left(\frac{999}{1000}\right)^{9990}$

- SUBWAYS** Fiona uses the subway to commute to work. During the morning commute, the trains run frequently. There is a 1 in 8 chance that she will find a train waiting for her as soon as she gets to the platform. Over the course of a five-day work week, what is the probability that she found a train waiting for her at least twice? Round your answer to the nearest thousandth.
0.121

- SOCCER** The boys varsity soccer team at Lincoln High School has a 75% probability of winning each of its 17 games this season. What is the probability that the team will win at least 13 games this season? Round your answer to the nearest thousandth.
0.574

CHESS For Exercises 6-8, use the following information.

Gary and Howard play chess. Gary's chess rating is 2050 and Howard's chess rating is 1948. This means that whenever they play, Gary has a 64% chance of defeating Howard. One day, Gary and Howard play three games against each other. Round your answers to the nearest thousandth.

- What is the probability that Gary will win all three of the matches?
0.262
- What is the probability that Gary will win at least two of the three matches?
0.705
- What is the probability that Gary will win only one of the matches?
0.249

NAME _____ DATE _____ PERIOD _____

12-9 Enrichment

Multinomial Experiments

A multinomial is a generalization of a binomial. For example, $(a + b + c)^2$ is a multinomial. One way to determine the coefficients is by direct multiplication using the distributive property. Take each term in the first factor and multiply by each term $(a, b, \text{ and } c)$ in the second factor, then combine like terms. (Notice that the sum of the exponents is always equal to two.)

$$\begin{aligned} (a + b + c)^2 &= (a + b + c)(a + b + c) = a(a + b + c) + b(a + b + c) + c(a + b + c) \\ &= a^2 + ab + ac + ab + b^2 + bc + ac + bc + c^2 \\ &= a^2 + b^2 + c^2 + 2ab + 2ac + 2bc \end{aligned}$$

Underlying this expansion is the notion of a *partition* into categories. The example partitions two 'items' among three categories. In this case the categories are the variables $a, b, \text{ and } c$ and the items are exponents. For example the partition, $\{1, 0, 1\}$ represents the term ac in the expansion, which could also be written as $a^1b^0c^1$, whereas the partition $\{0, 2, 0\}$ represents the b^2 term. The coefficients of each term can be computed by the formula:

$$\frac{n!}{n_1! \cdot n_2! \cdot n_3! \cdots n_k!}, \text{ where } n \text{ is the exponent and } n_1 + n_2 + n_3 + \cdots + n_k = n. \text{ Recall, } 0! = 1.$$

Term	Partition	Coefficient
a^2	$\{2, 0, 0\}$	$\frac{2!}{2! \cdot 0! \cdot 0!} = 1$
b^2	$\{0, 2, 0\}$	$\frac{2!}{0! \cdot 2! \cdot 0!} = 1$
c^2	$\{0, 0, 2\}$	$\frac{2!}{0! \cdot 0! \cdot 2!} = 1$
ab	$\{1, 1, 0\}$	$\frac{2!}{1! \cdot 1! \cdot 0!} = 2$
ac	$\{1, 0, 1\}$	$\frac{2!}{1! \cdot 0! \cdot 1!} = 2$
bc	$\{0, 1, 1\}$	$\frac{2!}{0! \cdot 1! \cdot 1!} = 2$

- Determine the all the *partitions* of $(x + y + z)^3$.
 $\{3, 0, 0\}, \{0, 3, 0\}, \{0, 0, 3\}, \{2, 0, 1\}, \{2, 1, 0\}, \{1, 2, 0\}, \{1, 0, 2\}, \{0, 1, 2\}, \{0, 2, 1\}, \{1, 1, 1\}$
- Determine the coefficients in the expansion of $(x + y + z)^3$ associated with each partition.
 $\{3, 0, 0\} \rightarrow 1, \{0, 3, 0\} \rightarrow 1, \{0, 0, 3\} \rightarrow 1, \{2, 0, 1\} \rightarrow 3, \{2, 1, 0\} \rightarrow 3, \{1, 2, 0\} \rightarrow 3, \{1, 0, 2\} \rightarrow 3, \{0, 1, 2\} \rightarrow 3, \{0, 2, 1\} \rightarrow 3, \{1, 1, 1\} \rightarrow 6$
- How can you build and interpret a trinomial distribution?
The trinomial distribution is interpreted as independent events with 3 outcomes whose individual probabilities are for example $x = 0.25, y = 0.45, \text{ and } z = 0.30$.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____ DATE _____ PERIOD _____

12-10 Study Guide and Intervention

Sampling and Error

Bias A sample of size n is random (or unbiased) when every possible sample of size n has an equal chance of being selected. If a sample is biased, then information obtained from it may not be reliable.

Example To find out how people in the U.S. feel about mass transit, people at a commuter train station are asked their opinion. Does this situation represent a random sample?

No; the sample includes only people who actually use a mass-transit facility. The sample does not include people who ride bikes, drive cars, or walk.

Exercises

Determine whether each situation would produce a random sample. Write *yes* or *no* and explain your answer.

- asking people in Phoenix, Arizona, about rainfall to determine the average rainfall for the United States **No; it rains less in Phoenix than most places in the U.S.**
- obtaining the names of tree types in North America by surveying all of the U.S. National Forests **Yes; there are National Forests in about every state in the U.S.**
- surveying every tenth person who enters the mall to find out about music preferences in that part of the country **Yes; mall customers should be fairly representative in terms of music tastes.**
- interviewing country club members to determine the average number of televisions per household in the community **No; country club members would tend to be more affluent and thus not a representative sample of the community.**
- surveying all students whose ID numbers end in 4 about their grades and career counseling needs **Yes; ID numbers are probably assigned alphabetically or by some other method not connected to students' grades or counseling needs.**
- surveying parents at a day care facility about their preferences for brands of baby food for a marketing campaign **Yes; choice of a daycare facility would probably not influence baby food preferences.**
- asking people in a library about the number of magazines to which they subscribe in order to describe the reading habits of a town **No; library visitors tend to read more than most citizens.**

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____ DATE _____ PERIOD _____

12-10 Lesson Reading Guide

Sampling and Error

Get Ready for the Lesson

Read the introduction to Lesson 12-10 in your textbook.

Do you think the results of the survey show that more mothers spend \$249 or less than \$250–\$349? If there is not enough information given to determine this, list at least two questions you would ask about the survey that would help you determine the significance of the survey. **Sample answer: There is not enough information to tell. 1. How many people were surveyed? 2. How was the sample for the survey selected? 3. What is the margin of error for this survey?**

Read the Lesson

- Determine whether each situation would produce a random sample. Write *yes* or *no* and explain your answer.
 - asking all the customers at five restaurants on the same evening how many times a month they eat dinner in restaurants to determine how often the average American eats dinner in a restaurant **No; people surveyed at a restaurant might be likely to eat dinner in restaurants more often than other people.**
 - putting the names of all seniors at your high school in a hat and then drawing 20 names for a survey to find out where seniors would like to hold their prom **Yes; every senior would have an equal chance of being chosen for the survey.**
- A survey determined that 58% of registered voters in the United States support increased federal spending for education. The margin of error for this survey is 4%. Explain in your own words what this tells you about the actual percentage of registered voters who support increased spending for education. **Sample answer: There is a 95% chance that the actual percentage of voters supporting increased federal spending for education is between 54% and 62%.**

Remember What You Learned

- The formula for margin of sampling error may be tricky to remember. A good way to start is to think about the variables that must be included in the formula. What are these variables, and what do they represent? What is an easy way to remember which variable goes in the denominator in the formula? **Sample answer: p is the probability of a certain response and n is the sample size. The larger the sample size, the smaller the margin of error, so n must go in the denominator since dividing by a larger number gives a smaller number. The square root of a smaller number is a smaller number, and twice the square root of a smaller number is a smaller number.**

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____ DATE _____ PERIOD _____

12-10 Study Guide and Intervention *(continued)*

Sampling and Error

Margin of Error The margin of sampling error gives a limit on the difference between how a sample responds and how the total population would respond.

Margin of Error If the percent of people in a sample responding in a certain way is p and the size of the sample is n , then 95% of the time, the percent of the population responding in that same way will be between $p - ME$ and $p + ME$, where $ME = 2\sqrt{\frac{p(1-p)}{n}}$.

Example 1 In a survey of 4500 randomly selected voters, 62% favored candidate A. What is the margin of error?

$$ME = 2\sqrt{\frac{p(1-p)}{n}}$$

Formula for margin of sampling error

$$= 2\sqrt{\frac{0.62 \cdot (1 - 0.62)}{4500}}$$

$$p = 62\% \text{ or } 0.62, n = 4500$$

Use a calculator.

$$\approx 0.01447$$

The margin of error is about 1%. This means that there is a 95% chance that the percent of voters favoring candidate A is between $62 - 1$ or 61% and $62 + 1$ or 63%.

Example 2 The CD that 32% of teenagers surveyed plan to buy next is the latest from the popular new group BFA. If the margin of error of the survey is 2%, how many teenagers were surveyed?

$$ME = 2\sqrt{\frac{p(1-p)}{n}}$$

Formula for margin of sampling error

$$0.02 = 2\sqrt{\frac{0.32 \cdot (1 - 0.32)}{n}}$$

$$ME = 0.02, p = 0.32$$

$$0.01 = \sqrt{\frac{0.32(0.68)}{n}}$$

Divide each side by 2.

$$0.0001 = \frac{0.32(0.68)}{n}$$

Square each side.

$$n = \frac{0.32(0.68)}{0.0001}$$

Multiply by n and divide by 0.0001

$$n = 2176$$

2176 teenagers were surveyed.

Exercises

Find the margin of sampling error to the nearest percent.

- $p = 45\%$, $n = 350$
about 5%
- $p = 12\%$, $n = 1500$
about 2%
- $p = 86\%$, $n = 600$
about 3%

- A study of 50,000 drivers in Indiana, Illinois, and Ohio showed that 68% preferred a speed limit of 75 mph over 65 mph on highways and country roads. What was the margin of sampling error to the nearest tenth of a percent? **about 0.4%**

Chapter 12

Glencoe Algebra 2

72

NAME _____ DATE _____ PERIOD _____

12-10 Skills Practice

Sampling and Error

Determine whether each situation would produce a random sample. Write yes or no and explain your answer.

- calling households at 3:30 P.M. on Tuesday to determine a political candidate's support
No; since most registered voters are likely to be at work at this time, this sample would not be representative of all registered voters.
- polling customers as they exit a sporting goods store about their attitudes about exercise
No; these customers are likely to value exercise more than those who do not shop at sporting goods stores, who are not represented in this survey.
- recording the number of sit-ups performed by 15-year-old girls in the high schools of a large school district to determine the fitness of all high-school girls in the district
No; 15-year-old girls may not have the same abilities as 18-year-old seniors, for example.
- selecting two of a city's 20 apartment buildings for a survey to determine the desire of apartment dwellers in the city to own a home
No; the residents of the two buildings selected might, for example, have nicer apartments or be in a nicer area of town, and thus would not well represent the desires of people in other buildings.
- In a large school district, the superintendent of schools interviews two teachers at random from each school to determine whether teachers in the district think students are assigned too much or too little homework.
Yes; since a cross section of teachers from all levels was selected at random, the sample should well represent the population of teachers in the district.
- For seven consecutive days, one hour each in the morning, afternoon, and evening, every tenth customer who enters a mall is asked to choose her or his favorite store.
Yes; because the sample is chosen over the course of a whole week, during hours when different consumer groups shop, and because the selection is systematic, the sample should well represent the general population that shops at the mall stores.

Find the margin of sampling error to the nearest percent.

- $p = 85\%$, $n = 100$
about 7%
- $p = 78\%$, $n = 100$
about 8%
- $p = 15\%$, $n = 100$
about 7%
- $p = 37\%$, $n = 500$
about 4%
- $p = 12\%$, $n = 500$
about 3%
- $p = 93\%$, $n = 500$
about 2%
- $p = 23\%$, $n = 1000$
about 3%
- $p = 56\%$, $n = 1000$
about 3%

- HEALTH** In a recent poll of cigarette smokers, 67% of those surveyed said they had tried to quit smoking within the last year. The margin of error was 3%. About how many people were surveyed? **about 983**

Chapter 12

Glencoe Algebra 2

73

NAME _____

DATE _____

PERIOD _____

12-10 Practice

Sampling and Error

Determine whether each situation would produce a random sample. Write *yes* or *no* and explain your answer.

- calling every twentieth registered voter to determine whether people own or rent their homes in your community **No; registered voters may be more likely to be homeowners, causing the survey to underrepresent renters.**
- predicting local election results by polling people in every twentieth residence in all the different neighborhoods of your community **Yes; since all neighborhoods are represented proportionally, the views of the community should as a whole should be well represented.**
- to find out why not many students are using the library, a school's librarian gives a questionnaire to every tenth student entering the library **No; she is polling only the students who are coming to the library, and will obtain no input from those who aren't using the library.**
- testing overall performance of tires on interstate highways only **No; for overall performance, tires should be tested on many kinds of surfaces, and under many types of conditions.**
- selecting every 50th hamburger from a fast-food restaurant chain and determining its fat content to assess the fat content of hamburgers served in fast-food restaurant chains throughout the country **No; the selected hamburgers are a random sample of the hamburgers served in one chain, and may represent the fat content for that chain, but will not necessarily represent the fat content of hamburgers served in other fast-food restaurant chains.**
- assigning all shift workers in a manufacturing plant a unique identification number; and then placing the numbers in a hat and drawing 30 at random to determine the annual average salary of the workers **Yes; because the numbers are randomly chosen from among all shift workers, all workers have the same chance of being selected.**

Find the margin of sampling error to the nearest percent.

- $p = 26\%$, $n = 100$ **about 9%**
- $p = 55\%$, $n = 100$ **about 10%**
- $p = 14\%$, $n = 500$ **about 3%**
- $p = 34\%$, $n = 1000$ **about 3%**
- $p = 49\%$, $n = 1500$ **about 3%**
- $p = 75\%$, $n = 500$ **about 4%**
- $p = 21\%$, $n = 1000$ **about 3%**
- $p = 65\%$, $n = 1500$ **about 2%**
- COMPUTING** According to a poll of 500 teenagers, 43% said that they use a personal computer at home. What is the margin of sampling error? **about 4%**
- TRUST** A survey of 605 people, ages 13–33, shows that 68% trust their parents more than their best friends to tell them the truth. What is the margin of sampling error? **about 4%**
- PRODUCTIVITY** A study by the University of Illinois in 1995 showed an increase in productivity by 10% of the employees who wore headsets and listened to music of their choice while they were working. The margin of sampling error for the study was about 7%. How many employees participated in the study? **about 73**

Chapter 12

74

Glencoe Algebra 2

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

NAME _____

DATE _____

PERIOD _____

12-10 Word Problem Practice

Sampling and Error

- COMICS** Isaac would like to know if people prefer reading comic books or novels. He decides to wait outside of a bookstore and ask people exiting whether they purchased comics or novels. Discuss whether this method of acquiring data would produce a biased or unbiased sample.
Sample answer: The results are likely to be biased because the customers of a bookstore may be more likely to purchase a novel over a comic book since bookstores do not generally offer a wide selection of comic books compared to novels.
- PARKING** A town wants to find out if people are happy with a proposal to tear down a section of a park and replace it with a parking lot. The town council decides to conduct a random survey of the town's citizens. They send a person to the location in the park where the proposed parking lot will be and have that person ask all passersby whether they would like to see a parking lot built at the location. Discuss whether or not this would produce a random sample.
it would not. Only people who use the park would be surveyed.
- PROMS** A poll asked 50 random seniors at a high school whether they would like to have the senior prom at a nearby hotel or at a local convention hall. Sixteen students responded that they would prefer the hotel. What is the margin of sampling error? Round your answer to the nearest percent.
13%
- AIRPORTS** In a large city, a random survey found that 18% of the city's population want a new runway built at the city airport. The survey had a margin of error of 5%. About how many people were surveyed?
about 236
- INTERNET USE For Exercises 5-7, use the following information.**
Two surveys were conducted to find out if people think that Americans are becoming more knowledgeable about the Internet. One survey polled 500 people and found that 395 of them felt that Americans are becoming more Internet savvy. A second survey concluded that 79% of those polled think that Americans are becoming more Internet savvy with a margin of error of 2%.
5. What was the margin of error for the first survey? Round your answer to the nearest percent.
4%
6. About how many people were polled in the second survey?
1659
7. Based on the results of the second survey, between what two percentages would you estimate is the true percentage of people who think that Americans are more Internet savvy, with 95% confidence?
77% and 81%

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Copyright © Glencoe/McGraw-Hill, a division of The McGraw-Hill Companies, Inc.

Chapter 12

75

Glencoe Algebra 2